

(12)

(21) **2 415 010**

(22) **04.07.2001**

(51) Int. Cl.<sup>7</sup>:

**C07D 235/06**, A61P 35/00,  
C07D 471/04, C07D 401/06,  
C07D 405/06, C07D 409/06,  
C07D 401/12, C07D 403/12,  
C07D 405/12, C07D 409/12,  
C07D 413/12, C07D 417/12,  
C07D 401/14, C07D 405/14,  
C07D 409/14, A61P 5/24,  
A61K 31/4184, A61K 31/422,  
A61K 31/427, A61K 31/428,  
A61K 31/4439, A61K 31/454,  
A61K 31/4725, A61K 31/506,  
A61K 31/5377, C07D 277/64

(85) **06.01.2003**

(86) **PCT/JP01/05813**

(87) **WO02/002533**

(30) 2000-204425 JP 05.07.2000  
2001-153372 JP 23.05.2001

(71) YAMANOUCI PHARMACEUTICAL CO. LTD.,  
3-11, Nihonbashi-Honcho 2-chome  
Chuo-ku  
103-8411, TOKYO, XX (JP).

KUSAYAMA, TOSHIYUKI (JP).  
MORITOMO, HIROYUKI (JP).  
TOYOSHIMA, AKIRA (JP).  
HIRANO, MASAOKI (JP).  
KAWAMINAMI, EIJI (JP).  
SEKI, NORIO (JP).  
OKADA, MINORU (JP).  
WAKAYAMA, RYUTARO (JP).

(72)

(74)

**DENNISON ASSOCIATES**

(54) DERIVES DE PROPANE-1,3-DIONE  
(54) PROPANE-1,3-DIONE DERIVATIVES



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2415010 A1 2002/01/10

(21) **2 415 010**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2001/07/04  
(87) **Date publication PCT/PCT Publication Date:** 2002/01/10  
(85) **Entrée phase nationale/National Entry:** 2003/01/06  
(86) **N° demande PCT/PCT Application No.:** JP 2001/005813  
(87) **N° publication PCT/PCT Publication No.:** 2002/002533  
(30) **Priorités/Priorities:** 2000/07/05 (2000-204425) JP;  
2001/05/23 (2001-153372) JP


(51) **Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup>** C07D 235/06, A61K 31/5377,  
A61P 35/00, A61K 31/506, A61K 31/4725, A61K 31/454,  
A61K 31/4439, A61K 31/428, A61K 31/427,  
A61K 31/422, A61K 31/4184, A61P 5/24, ...

(71) **Demandeur/Applicant:**  
YAMANOUCHI PHARMACEUTICAL CO. LTD., JP

(72) **Inventeurs/Inventors:**  
HIRANO, MASAOKI, JP;  
KAWAMINAMI, EIJI, JP;  
TOYOSHIMA, AKIRA, JP;  
MORITOMO, HIROYUKI, JP;  
SEKI, NORIO, JP;  
...

(74) **Agent:** DENNISON ASSOCIATES

(54) **Titre :** DERIVES DE PROPANE-1,3-DIONE  
(54) **Title:** PROPANE-1,3-DIONE DERIVATIVES



(57) **Abrégé/Abstract:**

Medicinal compositions, in particular, GnRH receptor antagonists comprising propane-1,3-dione derivatives represented by the following general formula (1) as the active ingredient; and novel propane-1,3-dione derivatives having a GnRH antagonism: general formula (1).

Canada

<http://opic.gc.ca> · Ottawa-Hull K1A 0C9 · <http://cipo.gc.ca>

OPIC · CIPO 191

OPIC



CIPO

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> (suite/continued) C07D 409/14, C07D 405/14, C07D 401/14, C07D 417/12, C07D 413/12, C07D 409/12, C07D 405/12, C07D 277/64, C07D 401/06, C07D 471/04, C07D 403/12, C07D 401/12, C07D 409/06, C07D 405/06

(72) Inventeurs(suite)/Inventors(continued): WAKAYAMA, RYUTARO, JP; OKADA, MINORU, JP; KUSAYAMA, TOSHIYUKI, JP

**Abstract**

Provided is a pharmaceutical composition containing a propane-1,3-dione derivative as the active ingredient, particularly a GnRH receptor antagonist. Also, provided is a  
5 propane-1,3-dione derivative having a GnRH antagonistic effect.

## Description

### PROPANE-1,3-DIONE DERIVATIVE

#### 5 Technical Field

The present invention relates to a pharmaceutical composition containing a propane-1,3-dione derivative or a pharmaceutically acceptable salt thereof as the active ingredient and a novel propane-1,3-dione derivative.

10


#### Background of the Invention

It is known that hypothalamic hormone or pituitary hormone takes part in a control system of secretion of peripheral hormones. In general, the secretion of anterior  
15 pituitary hormone is regulated by a secretion stimulating hormone or a secretion suppressing hormone secreted from an upper center, hypothalamus, a peripheral hormone secreted from the target organs them.

Gonadotropin releasing hormone (hereinafter,  
20 abbreviated as GnRH; also, GnRH is called as luteinizing hormone releasing hormone; LHRH) is known as a hormone which controls the secretion of sex hormones at the highest position, and regulates the secretion of luteinizing hormone (hereinafter, abbreviated as LH), follicle stimulating hormone  
25 (hereinafter, abbreviated as FSH), and sex hormones in the gonads through the intermediary of the receptor (hereinafter, abbreviated as GnRH receptor) which is considered to be

present in anterior pituitary (Horumon to Rinsyo (Hormones and Clinical), 46, 46-57 (1998)). A specific and selective antagonist against the GnRH receptor is expected to be a drug for preventing and treating sex hormone-dependent diseases since it regulates the action of GnRH and controls the secretion of lower LH, FSH and sex hormones (Horumon to Rinsyo (Hormones and Clinical) (1998), *ibid.*).

As compounds having a GnRH receptor antagonistic property, peptide compounds such as linear peptides, cyclic hexapeptide derivatives and bicyclic peptide derivatives which are derivatives of GnRH have been known. Also, as non-peptide compounds having the property, the following aminobenzimidazole derivatives (Japanese Patent Laid-Open No. 95767/2000), thienopyrimidine derivatives (WO 95/28405), or the like has been reported.



(refer to the above publications for the symbols in the formulae)

On the other hand, the known propane-1,3-dione derivatives having a benzimidazole, benzothiazole, or benzoxazole skeleton described in the following Table 1 have been reported as reagents for use as photosensitizer or the like (EP-A-135348, EP-A-631177, EP-A-368327, EP-A-332044, WO


94/01415, USP 4062686, USP 4119466, Collect. Czech. Chem. Commun. (1971), 36(1), 150-63, Zh. Nauch. Prikl. Fotogr. Kinematogr. (1971), 16(4), 282-8, Collect. Czech. Chem. Commun. (1978), 43(3), 739-45, Collect. Czech. Chem. Commun. (1979), 44(5), 1540-51, and Collect. Czech. Chem. Commun. (1973), 38(12), 3616-22, but pharmaceutical actions, particularly a GnRH receptor antagonistic action have not been disclosed.

10 Disclosure of the Invention

As a result of the intensive studies on non-peptide compounds having an excellent GnRH receptor antagonistic action, the present inventors have found that 2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-1,3-diphenylpropane-1,3-dione derivatives are useful. Furthermore, the inventors have developed various compounds based on the findings and have found that propane-1,3-dione derivatives represented by the following general formula (I) have an excellent GnRH receptor antagonistic action. Accordingly, they have accomplished the invention. Among the compounds of the invention, there are confirmed some compounds having a GnRH receptor-binding inhibitory activity equal to that of a peptide antagonist Cetrorelix which is commercialized at present. Thus, the invention relates to extremely useful compounds as non-peptide compounds.

Namely, the invention relates to the following:

That is, a pharmaceutical composition comprising a propane-1,3-dione derivative represented by the general formula (I):



5

(R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup>: the same or different, H, NO<sub>2</sub>, CN, Halo, a hydrocarbon group which may be substituted, a heterocycle which may be substituted, a hydroxy which may be substituted, a carboxy which may be substituted, an acyl-O- which may be substituted, an acyl which may be substituted, a substituent - S(O)<sub>n<sub>101</sub></sub>- (n<sub>101</sub>: an integer of 0 to 2, the same shall apply hereinafter), H-S(O)<sub>n<sub>101</sub></sub>-, a carbamoyl which may be substituted, a sulfamoyl which may be substituted, or an amino which may be substituted, and two adjacent groups selected from the group of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> may be combined to form an aryl or a cycloalkenyl;

R<sup>5</sup> and R<sup>6</sup>: the same or different, H, Halo, a hydrocarbon group which may be substituted or an amino which may be substituted;

X<sup>1</sup> and X<sup>2</sup>: the same or different, N, S or O atom;

A and B: the same or different, an aryl which may be substituted or a heterocycle which may be substituted;

Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup> and Z<sup>4</sup>: C or N;

provided that 1) when X<sup>1</sup> and X<sup>2</sup> each is S or O, one or both of the corresponding R<sup>5</sup> and R<sup>6</sup> are absent; 2) when one to four of










































$Z^1$ ,  $Z^2$ ,  $Z^3$  and/or  $Z^4$  are N, the corresponding  $R^1$ ,  $R^2$ ,  $R^3$  and/or  $R^4$  are absent)

or a pharmaceutically acceptable salt thereof as the active ingredient, preferably, the pharmaceutical composition which  
5 is a gonadotropin releasing hormone receptor antagonist, more preferably, the pharmaceutical composition comprising a propane-1,3-dione derivative wherein at least any one of  $X^1$  and  $X^2$  in the general formula (I) is N or the propane-1,3-dione derivative wherein  $X^1$  and  $X^2$  are N at the same time, or a  
10 pharmaceutically acceptable salt thereof as the active ingredient.

As another embodiment, the invention relates to a propane-1,3-dione derivative in the general formula (I) or a pharmaceutically acceptable salt thereof, provided that the  
15 compounds 1 to 39 shown in the following Table 1 are excluded, wherein the symbol Ph means phenyl, Me means methyl, Et means ethyl, or tBu means tert-butyl.

Table 1

1. 	14. 	27. 
2. 	15. 	28. 
3. 	16. 	29. 
4. 	17. 	30. 
5. 	18. 	31. 
6. 	19. 	32. 
7. 	20. 	33. 
8. 	21. 	34. 
9. 	22. 	35. 
10. 	23. 	36. 
11. 	24. 	37. 
12. 	25. 	38. 
13. 	26. 	39. 

Preferably, it is the propane-1,3-dione derivative wherein at least any one of  $X^1$  and  $X^2$  in the general formula (I) is N or the propane-1,3-dione derivative wherein  $X^1$  and  $X^2$  in the general formula (I) are N at the same time or a pharmaceutically acceptable salt thereof. In addition, as the other embodiment, it is the propane-1,3-dione derivative wherein  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  is H, an amino which may be substituted or a hydroxy which may be substituted, or a pharmaceutically acceptable salt thereof.

#### Best Mode for Carrying Out the Invention

The invention is further described in detail.

As the "Halo", fluorine, chlorine, bromine, or iodine atoms can be mentioned

The "hydrocarbon group" means a group composed of  $C_{1-15}$  carbons and hydrogens and is any form of linear or branched, monocyclic or fused polycyclic, and/or saturated or unsaturated ones, and preferably means an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, or aryl-alkyl.

The "alkyl" means a linear or branched saturated hydrocarbon group, and is preferably a  $C_{1-10}$  alkyl, more preferably a  $C_{1-6}$  alkyl. Specifically, it is methyl, ethyl, isopropyl, decyl, or the like. The "alkenyl" means a linear or branched hydrocarbon group having at least one or more double bonds, and is preferably a  $C_{2-10}$  alkenyl. Specifically, it is vinyl, propenyl, allyl, isopropenyl, hexenyl, or the like. The "alkynyl" means a linear or branched hydrocarbon

group having at least one or more triple bonds, and is preferably a C<sub>2-10</sub> alkynyl. Specifically, it is ethynyl, propynyl, butynyl, or the like. The "cycloalkyl" means a monocyclic saturated hydrocarbon ring, and is preferably a "C<sub>3-8</sub> cycloalkyl". Specifically, it is cyclopropyl, cyclopentyl, cyclohexyl, or the like. The "cycloalkenyl" means a monocyclic unsaturated hydrocarbon ring, and is preferably a "C<sub>3-8</sub> cycloalkenyl". Specifically, it is cyclopentenyl, cyclohexenyl, or the like. The "aryl" means an aromatic hydrocarbon ring, and is preferably C<sub>6-14</sub> aryl. Specifically, it is phenyl, naphthyl, 5,6,7,8-tetrahydro-naphthyl, indenyl, anthryl, fluorenyl, or the like.

The "heterocyclic group" means a five- or six-membered, monocyclic or bicyclic, saturated or unsaturated ring containing from 1 to 4 heteroatoms selected from N, S and O. The unsaturated ring includes aromatic rings (heteroaryls) and non-aromatic rings. As the monocyclic groups, there may be mentioned pyrrolidinyl, pyrazolidinyl, dioxanyl, piperadinyl, piperidinyl, morpholino, trithianyl, dioxolanyl, furyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, oxazolyl, pyridyl, pyrazinyl, pyrimidyl, triazolyl, thiadiazolyl, pyridazinyl, triazinyl, oxadiazolyl, or the like. As the bicyclic groups, there may be mentioned indolinyl, 3,4-methylenedioxyphenyl, 3,4-ethylenedioxyphenyl, benzofuranyl, benzothienyl, benzothiadiazolyl, benzothiazolyl, benzimidazolyl, indolyl, quinolyl, isoquinolyl, quinoxalinyl, or the like. It is preferably a five- or six-membered

monocyclic heteroaryl, more preferably furyl, thienyl, imidazolyl, thiazolyl, or pyridyl.

The "acyl" includes HCO-, a C<sub>1-15</sub> hydrocarbon group-CO-, a heterocyclic group-CO-, a heterocyclic group-alkyl-CO-, a heterocyclic group-alkenyl-CO-, a heterocyclic group-alkynyl-CO-, a C<sub>1-15</sub> hydrocarbon group-CS-, a heterocyclic group-CS-, a heterocyclic group-alkyl-CS-, a heterocyclic group-alkenyl-CS-, or a heterocyclic group-alkynyl-CS-. It is preferably a C<sub>1-15</sub> hydrocarbon group-CO- or a heterocyclic group-CO-, specifically HCO-, acetyl, propionyl, 2-methylbut-2-en-2-oyl, benzoyl, nicotinoyl, thenoyl, pyrrolidinylcarbonyl, piperidinylcarbonyl, or the like.

The "Halo hydrocarbon group" includes, for example, a Halo C<sub>1-10</sub> alkyl and a Halo C<sub>6-14</sub> aryl and specifically, chloromethyl, trifluoromethyl, fluorophenyl, difluorophenyl, trifluorophenyl, or the like.

The "heterocycle-ylidene" is a group wherein two bonding hands are present from the identical carbon atom in a heterocycle, and examples thereof include 3-methyl-4-oxo-2-thioxothiazolidin-5-ylidene and the like.

The "heterocyclic group-C<sub>1-10</sub> alkylidene" includes pyridylmethylenidene and the like.

The substituents in the hydrocarbon group which may be substituted include specifically the substituents in the following a group.

The substituents in the heterocycle which may be substituted, the hydroxy which may be substituted, the carboxy

which may be substituted, the acyl-O- which may be substituted, the acyl which may be substituted, the substituent  $-S(O)_{n_{101}}-$ , the carbamoyl which may be substituted, and the sulfamoyl which may be substituted include specifically the substituents in the following b group.

The substituents in the amino which may be substituted include specifically the substituents in the following c group.

Moreover, the substituents in the aryl or heterocyclic group which may be substituted in A ring and B ring include the substituents in the following d group.

a group: OH, NO<sub>2</sub>, COOH, Halo, C<sub>6-14</sub> aryl, heterocyclic group, R<sup>101</sup><sub>3</sub>SiO- and/or R<sup>101</sup>-T<sup>101</sup>-

R<sup>101</sup>: (1) H, (2) C<sub>3-8</sub> cycloalkyl, (3) heterocyclic group, (4) C<sub>1-10</sub> alkyl which may be substituted by [OH, NO<sub>2</sub>, COOH, Halo, heterocyclic group, C<sub>1-10</sub> alkyl-CO-, C<sub>1-10</sub> alkyl-O-, C<sub>1-10</sub> alkyl-O-CO-, and/or (R<sup>102</sup>)<sub>n<sub>102</sub></sub>C<sub>6-14</sub> aryl],  
R<sup>102</sup>: H, Halo, NO<sub>2</sub>, OH, COOH, C<sub>1-10</sub> alkyl-O- or C<sub>1-10</sub> alkyl-O-CO-,

n<sub>102</sub>: an integer of 1 to 5,

(5) C<sub>6-14</sub> aryl which may be substituted by OH, CN, NO<sub>2</sub>, Halo, and/or C<sub>1-10</sub> alkyl-CONR<sup>103</sup>-,

R<sup>103</sup>: the same as or different from R<sup>101</sup>,

(a) H, (b) C<sub>3-8</sub> cycloalkyl, (c) heterocyclic group,

(d) C<sub>1-10</sub> alkyl which may be substituted by COOH, C<sub>1-10</sub> alkyl-O-CO-, (R<sup>104</sup>)<sub>n<sub>102</sub></sub>-C<sub>6-14</sub> aryl or (R<sup>104</sup>)<sub>n<sub>102</sub></sub>-heterocyclic group,

R<sup>104</sup>: H, OH, Halo or C<sub>1-10</sub> alkyl-O-, or

5 (e) C<sub>6-14</sub> aryl which may be substituted by OH, CN, NO<sub>2</sub>, Halo or C<sub>1-10</sub> alkyl-CONR<sup>105</sup>-,

R<sup>105</sup>: (a) H, (b) C<sub>3-8</sub> cycloalkyl, (c) heterocyclic group,

10 (d) C<sub>1-10</sub> alkyl which may be substituted by COOH, C<sub>1-10</sub> alkyl-O-CO-, C<sub>6-14</sub> aryl or heterocyclic group, or

(e) C<sub>6-14</sub> aryl which may be substituted by OH, CN, NO<sub>2</sub> or Halo,

T<sup>101</sup>: -O-, -CO-, -CO-O-, -O-CO-, -NR<sup>103</sup>-CO- or -NR<sup>103</sup>-, the same shall apply hereinafter,

15 b group: (1) H, (2) C<sub>3-8</sub> cycloalkyl, (3) C<sub>6-14</sub> aryl which may be substituted by C<sub>1-10</sub> alkyl-O-, (4) heterocyclic group, (5) C<sub>1-10</sub> alkyl which may be substituted by (OH, NO<sub>2</sub>, Halo, heterocyclic group, R<sup>101</sup>R<sup>103</sup>N, C<sub>1-10</sub> alkyl-O-, acyl or (R<sup>106</sup>)<sub>n<sub>102</sub></sub>-C<sub>6-14</sub> aryl),

R<sup>106</sup>: H, COOH, NO<sub>2</sub>, R<sup>101</sup>R<sup>103</sup>N, acyl-NR<sup>101</sup>- or C<sub>1-10</sub> alkyl-O-

20 CO-, the same shall apply hereinafter,

c group: (1) heterocyclic group which may be substituted by C<sub>1-10</sub> alkyl, Halo C<sub>1-10</sub> alkyl or C<sub>6-14</sub> aryl-C<sub>1-10</sub> alkyl, (2) C<sub>6-14</sub> aryl which may be substituted by cycloalkyl or R<sup>101</sup>R<sup>103</sup>N, (3) C<sub>1-10</sub> alkyl which may be substituted by R<sup>107</sup>,

25 R<sup>107</sup>: (a) C<sub>3-8</sub> cycloalkyl, (b) C<sub>3-8</sub> cycloalkenyl, (c) R<sup>108</sup>-O-,

$R^{108}$ : (i)  $C_{1-10}$  alkyl which may be substituted by  $C_{6-14}$  aryl, heterocyclic group or  $R^{101}R^{103}N$ , or (ii) aryl which may be substituted by  $C_{6-14}$  aryl or  $R^{101}R^{103}N$ ,

(d) acyl which may be substituted by  $NO_2$ , (e)  $(R^{109})_{n_{102}}-C_{6-14}$  aryl

$R^{109}$ : (i) H, (ii) OH, (iii) CN, (iv)  $NO_2$ , (v) COOH, (vi) Halo, (vii) oxo ( $=O$ ), (viii)  $R^{101}R^{103}N$ , (ix)  $C_{1-10}$  alkyl which may be substituted by  $R^{110}$ ,

$R^{110}$ : H, OH, COOH, Halo,  $C_{6-14}$  aryl,

heterocycle-ylidene which may be

substituted by ( $C_{1-10}$  alkyl, oxo or thioxo ( $=S$ )),  $C_{1-10}$  alkyl-O-,  $C_{1-10}$  alkyl-O-CO- or

acyl-O-, (x) acyl-O-, (xi)  $C_{6-14}$  aryl which may be substituted by Halo, (xii)

heterocyclic group which may be substituted by Halo,  $C_{1-10}$  alkyl or Halo  $C_{1-10}$  alkyl, and/or (xiii)  $R^{111}-T^{102}-$

$R^{111}$ ; (i) H, (ii)  $C_{3-8}$  cycloalkyl, (iii)

$R^{101}R^{103}N$ , (iv)  $C_{6-14}$  aryl which may be

substituted by Halo,  $C_{1-10}$  alkyl, Halo  $C_{1-10}$  alkyl or  $C_{6-14}$  aryl which may be  $C_{6-14}$  aryl,

or (v)  $C_{1-10}$  alkyl which may be substituted by Halo, COOH,  $C_{1-10}$  alkyl-O-,  $R^{101}R^{103}N$ ,  $C_{6-14}$  aryl, heterocyclic group, heterocycle-ylidene,  $C_{1-10}$  alkyl-O-CO- or acyl-O-



$T^{102}$ :  $-O-$ ,  $-CO-$ ,  $-NR^{101}-$ ,  $-O-CO-$ ,  $-CONR^{101}-$ ,  
 $-NR^{101}NR^{101}CO-$ ,  $-O-CONR^{101}-$ ,  $-S(O)_{n_{101}}-$  or  
 $-S(O)_{n_{101}}NR^{101}-$  and/or  $R^{111b}NC(NR^{111b})NR^{101}-$ ,  
 $R^{111b}$ : H or  $C_{1-10}$  alkyl-O-CO-

5 (f)  $(R^{112})_{n_{102}}$ -heterocyclic group,

$R^{112}$ : oxo, oxide or a group the same as  $R^{109}$

(g)  $C_{1-10}$  alkyl-O-CO-

(4) heterocyclic group- $C_{1-10}$  alkylidene which may be substituted  
 by Halo, oxide,  $C_{1-10}$  alkyl,  $C_{1-10}$  alkyl-O- or  $C_{1-10}$  alkyl-O-CO-  
 10  $NR^{101}-$ ,

(5) acyl which may be substituted by  $R^{113}$

$R^{113}$ : OH, COOH, CN,  $NO_2$ , Halo,  $C_{6-14}$  aryl, heterocyclic  
 group,  $R^{101}R^{103}N$ ,  $C_{1-10}$  alkyl, Halo  $C_{1-10}$  alkyl,  $C_{1-10}$   
 alkyl-O-,  $C_{1-10}$  alkyl-O-CO-,  $C_{1-10}$  alkyl-O- $C_{6-14}$  aryl,  
 15 acyl,  $C_{1-10}$  alkyl-O-CO-,  $C_{1-10}$  alkyl- $C_{6-14}$  aryl, acyl-  
 $NR^{101}-$ , acyl- $NR^{101}-C_{6-14}$  aryl or  $C_{1-10}$  alkyl- $C_{6-14}$  aryl- $SO_2-$   
 $NR^{101}-$ ,

(6)  $R^{101}R^{103}NCO$

(7)  $R^{114}-S(O)_{n_{101}}-$

20  $R^{114}$ : (a) H, (b)  $C_{1-10}$  alkyl which may be substituted by OH,  
 $NO_2$ , Halo,  $R^{101}R^{103}N$ ,  $C_{1-10}$  alkyl-O-, acyl- $NR^{101}-$  or  $C_{6-14}$  aryl,  
 (c)  $C_{6-14}$  aryl which may be substituted by OH,  $NO_2$ , Halo,  
 $R^{101}R^{103}N$ ,  $C_{1-10}$  alkyl, Halo  $C_{1-10}$  alkyl,  $C_{1-10}$  alkyl-O-, acyl-  
 $NR^{101}-$  or  $C_{6-14}$  aryl, (d) heterocyclic group which may be  
 25 substituted by OH,  $NO_2$ , Halo,  $R^{101}R^{103}N$ ,  $C_{1-10}$  alkyl, Halo  
 $C_{1-10}$  alkyl,  $C_{1-10}$  alkyl-O-, acyl- $NR^{101}-$  or  $C_{6-14}$  aryl, or (e)  
 $R^{101}R^{103}N$ , and/or

(8)  $R^{115}-T^{103}-$

$R^{115}$ : (a)  $C_{1-10}$  alkyl which may be substituted by heterocyclic group, (b)  $C_{6-14}$  aryl which may be substituted by heterocyclic group or  $R^{101}R^{103}N$  or (c) heterocyclic group,

$T^{103}$ :  $-CO-NR^{101}-$ ,  $-NR^{101}-CO-$ ,  $-NR^{101}-CS-$ ,  $-O-CO-CO-$ ,  $-O-CO-$  or  $-CO-CO-$ , the same shall apply hereinafter;


d group: (1) CN, (2)  $NO_2$ , (3) Halo, (4) OH, (5) COOH, (6)  $C_{1-10}$  alkyl- $T^{104}-$  which may be substituted by (OH, Halo, heterocyclic group,  $C_{6-14}$  aryl which may be substituted by Halo,  $R^{101}R^{103}N$ ,  $R^{101}-CO-$ ,  $R^{101}-T^{101}-CO-$  or  $R^{101}-T^{101}-$ ),

$T^{104}$ : a bond,  $-O-$ ,  $-CO-O-$ ,  $-O-CO-$ ,

(7) acyl which may be substituted by  $R^{113}$ , (8) acyl-O- which may be substituted by  $R^{113}$ , (9)  $R^{116}R^{117}N$


$R^{116}$ ,  $R^{117}$ : the same or different, H or a substituent of c group, and/or (10)  $R^{116}R^{117}NCO$ , the same shall apply hereinafter.)

In the active ingredients of the invention or the compounds of the invention, geometrical isomers and tautomers may exist, for example, as shown below.



The invention includes separated or mixed forms of these isomers. In addition, depending on the kind of substituents, certain compounds of the invention may contain asymmetric atom(s) or axial asymmetry, and hence isomers based on the asymmetric carbon atom(s) or the like can exist. The invention includes mixed or separated forms of these optical isomers. Moreover, the invention also includes compounds labeled with a radioactive isotope.

In addition, among the compounds of the invention, there exist compounds wherein geometrical isomerism with regard to the double bond at 2-position of the propane can be mutually transformable as shown below through tautomerism as shown in the above.



Furthermore, the active ingredients of the invention or the compounds of the invention also forms acid addition salts or salts with a base in some cases depending on the kinds of substituents, and such salts are also included in the invention so far as they are pharmaceutically acceptable salts. Specifically, there are mentioned acid addition salts with an inorganic acid such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, nitric acid or phosphoric acid or with an organic acid such as formic acid, acetic acid, propionic acid, oxalic acid, malonic acid, succinic acid, fumaric acid, maleic acid, lactic acid, malic acid, tartaric acid, citric acid, methanesulfonic acid, ethanesulfonic acid, aspartic acid or glutamic acid, salts with an inorganic base such as sodium, potassium, magnesium, calcium or aluminum or with an organic base such as methylamine, ethylamine, ethanolamine, lysine or ornithine, ammonium salts, and the like. Various hydrates and solvates of the active ingredients of the invention or the compounds of the invention are also included in the invention. In addition, polymorphic substances thereof are also included.

Moreover, the active ingredients of the invention or the compounds of the invention also include all the compounds which are metabolized and converted in the living body, so-called prodrugs. The groups forming the prodrugs of the invention include groups described in Prog. Med., 5, 2157-2161 (1985) and "Iyakuhiin no Kaihatsu (Development of

Pharmaceuticals)", Vol. 7 (Hirokawa Shoten, 1990), Bunshi Sekkei (Molecular Design), pp. 163-198 or the like.

(Production methods)

5           The compounds of the invention and the pharmaceutically acceptable salts thereof can be produced by utilizing characteristics based on the fundamental skeleton or kind of substituents and applying various known synthetic methods.

10           At that time, depending on the kind of functional group, it may sometimes be effective from the viewpoint of production techniques to replace the functional group with an appropriate protective group (a group which can be easily converted into the functional group) at the stage of starting materials or synthetic intermediates. Examples of such functional groups include an amino group, a hydroxyl group, a carboxyl group and the like and examples of their protective groups include the protective groups which are described in "Protective Groups in Organic Synthesis (3rd edition)",  
15           written by Greene and Wuts, which may be optionally selected and used depending on the reaction conditions. In these methods, the reaction is carried out after introducing a protective group and then, if necessary, the protective group is removed to obtain the desired compound.

25           Moreover, when the active ingredients of the invention are known compounds, they can be easily available in

accordance with the above literatures (Collect. Czech. Chem. Commun. (1971), 36(1), 150-163 and so forth).

The following will describe representative synthetic methods of the compounds of the invention or intermediates thereof.

The symbols in the following sentences are as follows.

DMF: N,N-dimethylformamide; DMSO: dimethyl sulfoxide; THF: tetrahydrofuran; Tol: toluene; EtOAc: ethyl acetate; DCE: 1,2-dichloroethane; TEA: triethylamine; Diglyme: diethylene glycol dimethyl ether


#### First production method (acylation reaction)

The present production method is a usual acylation, which is specifically carried out by reacting an alkyl compound with an equivalent amount of acyl compound in a solvent inert to the reaction at a room temperature to an elevated temperature.



The solvent inert to the reaction includes aromatic hydrocarbon solvents such as benzene or toluene, ether solvents such as Diglyme, THF, 1,4-dioxane or 1,2-dimethoxyethane, Halo hydrocarbon solvents such as dichloromethane, chloroform or DCE, basic solvents such as TEA, pyridine, collidine, morpholine or 2,6-lutidine, and the like. These solvents are used solely or as a mixture of two or more of them. Optionally, an inorganic base such as sodium hydride may be added.

As an representative example, the compound of the invention is produced by reacting a methylimidazole compound (II) with an acyl compound (III) in a solvent inert to the reaction at a room temperature to an elevated temperature (Step i) to obtain an intermediate (IV) or the like and by adding an equivalent amount of a carboxylic acid (V) to the compound (IV) and heating them (Step ii).

In the production method, the reaction can be also effected by adding an equivalent amount of a carboxylic acid (V) or an equivalent amount of water after the first step without isolating the intermediate (IV) or the like and heating them as above. Moreover, an acid anhydride of the acyl compound (III) may be used instead of the compound.



(wherein the symbol  $L^1$  in the formula represents a leaving

group, the above  represents  ; the same shall apply hereinafter)

The leaving group  $L^1$  includes Halo, or an organic sulfonic acid residue such as methanesulfonyloxy or p-toluenesulfonyloxy.


#### Second production method

The production method is carried out by reacting an ester compound (VII) with an acyl compound (VIII) to obtain a diketone compound (IX), by reacting the compound (IX) with carbon disulfide and adding an alkyl halide to obtain a dithioacetal compound (X), and finally by reacting the compound (X) with an amine compound (XI).

The compound of the invention (XII) is obtained by reacting the ester compound (VII) with an equivalent amount of the acyl compound (VIII) in the presence of a base such as sodium hydride in a solvent inert to the reaction, such as THF, at a room temperature to an elevated temperature (Step i), reacting the resulting compound (IX) with carbon disulfide in the presence of an inorganic base such as  $KF/Al_2O_3$  or potassium carbonate or an organic base such as TEA at a temperature of cooled temperature to room temperature, preferably  $0^\circ C$  to room temperature, then adding an alkylating agent such as methyl iodide or 1,3-dibromopropane to effect alkylation reaction (Step ii), and finally reacting the resulting dithioacetal compound (X) with an equivalent amount



of the amine compound (XI) in a solvent inert to the reaction, such as ethanol or DMSO, at a room temperature to under heating with refluxing (Step iii).




(wherein  $R^7$  represents a  $C_{1-6}$  alkyl and  $X^1$  represents NH, O or S, and a dotted line represents possible formation of a ring through combination of the two alkyl groups)

### 10 Third production method

The production method is carried out by reacting the 2-methylimidazole compound (II) with the acyl compound (III) (Step i), obtaining an imidazole compound (XIV) in the presence of an organic base such as morpholine in a solvent inert to the reaction at a room temperature to an elevated temperature (Step ii), and acylating the compound (XIV) with an acyl compound (XV) (Step iii). Step i and Step iii are carried out in accordance with the acylation in the above First production method. The intervening intermediate (XIII) or the like may be isolated or may not be isolated.

15

20




#### Fourth production method (reduction reaction)


The reduction reaction is carried out according to well-known methods (COMPREHENSIVE ORGANIC SYNTHESIS 8 REDUCTION (Pergamon Press) (1991)). More preferably, it is carried out by (1) catalytic reduction under hydrogen atmosphere or in the presence of hydrogen donor such as ammonium formate using palladium (Pd), platinum (Pt), nickel (Ni), or the like in a solvent such as methanol, ethanol, chloroform, EtOAc or acetic acid at a room temperature to an elevated temperature, (2) using a metal such as Fe or SnCl<sub>2</sub> in the presence of an acid such as acetic acid, hydrochloric acid or the like, or using a reducing agent such as sodium hydrosulfite in an mixed solvent such as water and MeOH or THF at a room temperature to an elevated temperature, or (3) adding a reducing agent such as sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride in a solvent inert to the reaction, such as ethanol, at a temperature of ice-cooling to an elevated temperature.

As a representative example, there may be mentioned a reaction from a nitro compound (XVII) to an amine compound (XVIII) or a reaction from a ketone compound (XIX) to an alcohol compound (XX).

5



or



10

(the symbol  $\text{R}^8$  in the formulae is a hydrocarbon group which may be substituted or a heterocyclic group which may be substituted,  $m1$  or  $m2$  is the same or different and represents an integer of 0 to 5,  $m3$  represents an integer of 0 to 4, and they satisfies  $m1+m2+m3 \geq 1$ ; the same shall apply hereinafter)

15


#### Fifth production method

The reaction is carried out by stirring an amine compound and an equivalent amount of an aldehyde compound in the presence or absence of an acid such as p-toluenesulfonic acid in a solvent inert to the reaction, such as ethanol, benzene, THF or Tol at a room temperature to an elevated temperature to obtain an imine compound, and then subjecting

20


it to reduction reaction in accordance with the above fourth production method, preferably the reaction (1) or (3).

Alternatively, the reaction is carried out by mixing an amine compound and an equivalent amount of an aldehyde compound and adding a reducing agent in accordance with Fourth production method. The reducing agent may be added immediately after the mixing of the amine compound and the aldehyde compound or at an interval of some period of time. A ketone or 1-hydroxymethylbenzotriazole may be used instead of the aldehyde compound. As representative examples, there may be mentioned a reaction from an amine compound (XVIII) and an aldehyde compound (XXI) to an alkylamino compound (XXII) and a reaction from the amine compound (XVIII) to the alkylamino compound (XXII) via an imine compound (XXIII).



(wherein R<sup>9</sup> represents the following meaning.

R<sup>9</sup>: R<sup>10</sup>-T<sup>1</sup>-

20 R<sup>10</sup>: H; R<sup>107</sup>;  (R<sup>e</sup>, R<sup>f</sup>: the same or different, hydrogen atom, or a substituent of the above c group,

a dotted line: R<sup>e</sup> and R<sup>f</sup> may be combined to form the above heterocycle (the same shall apply hereinafter); or a C<sub>1-15</sub> hydrocarbon group which may have 1 to 5 functional groups selected from the group consisting of C<sub>1-10</sub> alkyl-CONH-, C<sub>1-10</sub> alkyl, C<sub>1-10</sub> alkyl-O-, and a carboxy which may be substituted by a substituent of the above b group,

T<sup>1</sup>: C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl or a single bond, the same shall apply hereinafter)


Sixth production method (amidation or sulfonamidation reaction)

The reaction is carried out according to usual methods. For example, it is carried out according to a method using a condensing reagent (dicyclohexylcarbodiimide, 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide, 1,1'-carbonyldiimidazole, or the like) or a mixed acid anhydride method using ethyl chloroformate, isobutyl chloroformate, or the like.

Moreover, it is possible that a carboxylic acid or a sulfonic acid is converted into a reactive derivative such as an acid halide with a halogenating agent such as thionyl chloride, oxalyl chloride or phosphorus oxychloride and then the derivative is reacted with an amine compound. The reaction is usually suitably carried out in a solvent inert to the reaction, such as THF, DMF, dichloromethane, chloroform, acetonitrile or EtOAc in the presence of, if necessary, an organic base such as TEA or an inorganic base such as

potassium carbonate under cooling (preferably -15 to 0°C) or at room temperature or under heating.

As a representative example, there may be mentioned a reaction from the amine compound (XVIII) and a carboxylic acid (XXIV) or a reactive derivative thereof or a sulfonic acid (XXVI) or a reactive derivative thereof to an amide compound (XXV) or a sulfonamide compound (XXVII).




(the symbol in the formulae represents the following meaning.  
 $R^{11}CO$ : an acyl which may be substituted by  $R^{113}$ , the same shall apply hereinafter)

#### Seventh production method


The production method is carried out by reacting a compound having a leaving group with an equivalent amount of an amine compound, a compound having a hydroxy (OH) group or a sulfonamide compound in a solvent inert to the reaction, such as THF, acetone, DMF, acetonitrile, dichloromethane, methanol or DMSO, under cooling or at a room temperature to an elevated temperature, or under refluxing. Optionally, an

inorganic base such as potassium carbonate or an organic base such as TEA may be added.

As representative examples, there may be mentioned an amination reaction from an alkyl compound having a leaving group L2 (XXVIII) and an amine compound (XXIX) to a compound of the invention (XXX) or an O-alkylation reaction from an alkyl compound having a leaving group L2 (XXXIV) and a hydroxyl compound (XXXIII) to a compound of the invention (XXXV).



(the symbols in the formulae are as follows. L2: the above L1 or a diazo group (N=N-), L3: chlorine (Cl) or bromine (Br),  
 15 n1, n3: the same or different, an integer of 0 to 1,  
 n2: an integer of 0 to 4,  
 provided that m1+m2+n1+n2+n3<sub>≥</sub>1 (the same shall apply hereinafter), or




(the symbol  $R^{12}$  in the formulae is a substituent of b group, the same shall apply hereinafter)

#### 5 Eighth production method

The present hydrolysis reaction is carried out in the presence of an inorganic base such as potassium carbonate, more preferably an organic base such as morpholine in a solvent inert to the reaction at a room temperature to under heating with refluxing.

As a representative example, there may be mentioned a hydrolysis reaction from a compound (XXXI) to a compound (XXXII).



#### Ninth production method

The production method is carried out by reacting an amine compound with an equivalent amount of an isocyanate compound or an isothiocyanate compound in a solvent inert to the reaction such as Tol, acetonitrile, chloroform or DMF at a temperature of 0°C to under refluxing.




The isocyanate compound is obtained by subjecting a carboxylic acid or a reactive derivative thereof (e.g., an acid chloride), which is a starting material of the isocyanate compound, to well-known rearrangement reaction (ADVANCED  
5 ORGANIC CHEMISTRY written by J. March (John Willy & Sons (1992))). The isothiocyanate compound is obtained by subjecting an amine compound, alkyl halide, diazonium salt or isocyanide, which is a starting material of the isocyanate compound, to a well-known reaction (ADVANCED ORGANIC CHEMISTRY  
10 written by J. March (John Willy & Sons (1992))).

The isocyanate compound or isothiocyanate compound is obtained by the above reaction, and the compound may be subjected to a urea-forming reaction or thiourea-forming reaction in situ, or the isocyanate compound or isothiocyanate  
15 compound may be subjected to a urea-forming reaction or thiourea-forming reaction after once isolated. Alternatively, the production method is carried out by reacting an amine compound with an equivalent amount of a carbodiimide compound instead of the isocyanate compound or isothiocyanate compound  
20 in a solvent inert to the reaction at a room temperature to an elevated temperature and then subjecting the product to a deprotection reaction. The carbodiimide compound is synthesized by a well-known reaction (Fieser and Fieser's Reagent for Organic Synthesis, Vol. 8 (Wiley) p. 96). The  
25 carbodiimide compound may be protected with an appropriate protective group. The protective group and deprotection

reaction are in accordance with the above "Protective Groups in Organic Synthesis (third edition)".

As representative examples, there may be mentioned a reaction from an amine compound (XXXVI) and an isocyanate compound or isothiocyanate compound to a urea compound (XXXVII) or thiourea compound (XXXVIII) or a reaction from the amine compound (XXXVI) and a carbodiimide compound (XXXIX) to a guanidine compound (XXXXI).




10

#### Tenth production method

The present oxidation reaction is carried out according to well-known methods (ADVANCED ORGANIC CHEMISTRY written by J. March (John Willy & Sons (1992))). It is preferably carried out in a solvent inert to the reaction such as dichloromethane, chloroform, or the like in the presence of

an oxidizing agent such as m-chloroperbenzoic acid (mcpba), hydrogen peroxide or tetrapropylammonium perruthenate (TPAP).

As representative examples, there may be mentioned a reaction from a sulfide compound (XXXXII) and an oxidizing agent to a sulfonyl compound (XXXXIII), a reaction of an alcohol compound to an aldehyde compound, or a reaction from a pyridylmethylamino compound to an N-oxidopyridylmethylideneamino compound.



It is noted that each reaction scheme described in the above production methods show a reaction of a representative compound. Therefore, when the same substituent is present in the compounds of the invention at a position other than the position in the reaction scheme, the compounds included in the scope of the invention can be easily produced by the substituent-modifying reaction using the above reaction scheme.


Moreover, when the starting compounds are novel, they may be obtained by the following production methods.


#### Production method 1

The production method is carried out by condensing an aldehyde compound or ketone compound with an equivalent amount of an active methylene compound in the presence of a base or

an acid catalyst at a room temperature to an elevated temperature.


Acetic acid is used as a solvent, a secondary amine such as piperidine is preferably employed as the base, and a  
 5 salt such as ammonium chloride or potassium fluoride or a Lewis acid such as  $\text{TiCl}_4$  is used as the acid catalyst.



10 (wherein  is a heterocycle having oxo and active methylene, which may be substituted by C<sub>1-10</sub> alkyl and/or thioxo)

#### Production method 2


The production method is carried out by reacting a  
 15 nitrobenzene compound having a leaving group  $\text{L}^1$  with an equivalent amount of ammonia in a solvent inert to the reaction, such as methanol, at a room temperature to an elevated temperature in a sealed tube.



#### Production method 3

The production method is effected by reacting an aldehyde compound or ketone compound with an equivalent amount


of a phosphorus ylide in a solvent inert to the reaction, such as DMF, at a temperature of 0°C to an elevated temperature. The phosphorus ylide may be prepared from a corresponding phosphonium salt and a base such as sodium hydride according to well-known methods (ADVANCED ORGANIC CHEMISTRY written by J. March (John Willy & Sons (1992))).



(wherein  $R^{13}$  is a heteroaryl or a hydrocarbon group which may be substituted)

#### Production method 4

The production method is carried out by reacting a 1,2-phenylenediamine compound with a trialkyl orthoacetate compound in a solvent inert to the reaction, such as ethanol, at a room temperature to under refluxing. As occasion demands, an acid catalyst such as hydrochloric acid may be added, or water may be removed from the reaction system by adding molecular sieves or using the Dean-Stark apparatus.




Alternatively, the production method is carried out by cyclization-condensation of an o-aminoacetanilide compound

obtainable by carrying out a reduction reaction in accordance with the Fourth production method in the presence or absence of an acid catalyst such as acetic acid or hydrochloric acid. At that time, the resulting o-aminoacetanilide compound may be isolated or may not be isolated.

#### Production method 5

The production method is carried out by reacting a halobenzene compound with a phenylborane compound in the presence of a palladium catalyst such as tris(dibenzylideneacetone)dipalladium or the like and a ligand such as tris-t-butylphosphine or the like and a base such as cesium carbonate in a solvent inert to the reaction, such as dioxane, at a room temperature to an elevated temperature (Angew. Chem. Int. Ed., 37, 3388 (1998)). The reaction is preferably carried out under an atmosphere of an inert gas such as nitrogen gas or argon.




#### Production method 6

The production method is carried out by reacting an amine compound, amide compound or imide compound with an equivalent amount of a bromine compound in a solvent inert to the reaction, such as a Halo hydrocarbon solvent such as tetrachloromethane or DCE or an aromatic hydrocarbon solvent

such as benzene at a temperature of 0°C to under refluxing. The bromine compound includes N-bromosuccinimide, bromine, tBuOBr, AcOBr, and the like. Optionally, a radical initiator such azobisisobutyronitrile (AIBN) may be added.

5



The compounds of the invention are isolated and purified as free compounds, pharmaceutically acceptable salts thereof, hydrates, solvates or polymorphic substances. The pharmaceutically acceptable salts of the compounds of the invention (I) can be also produced by subjecting the compound to a usual salt-forming reaction.

Isolation and purification are carried out by applying general chemical operations such as extraction, fractionating, recrystallization, various types of fractionating chromatography and the like.

Each form of isomers can be separated by selecting an appropriate starting compound or making use of physicochemical differences among isomers. For example, optical isomers can be made into stereochemically pure isomers by selecting appropriate starting compounds or a conventional optical resolution method (e.g., a method in which they are converted into diastereomer salts with a general optically active base or acid and then subjected to optical resolution).

In a similar manner to those described in the production methods, the Example compounds described in the following tables are obtained. Also, part of the compounds in the following tables were obtained.


5 In this connection, abbreviations herein indicate as follows.

Rex: Reference Example; Ex: Example; Str: structural formula; Dat: physicochemical properties; FA: FAB-MS (M+H)<sup>+</sup>; MS: found value on mass spectrometry; FN: FAB-MS (M-H)<sup>-</sup>; EI: EI-MS; N1: 10 NMR (DMSO-d<sub>6</sub>, TMS internal standard) characteristic peaks  $\delta$  ppm; N2: NMR (CDCl<sub>3</sub>, TMS internal standard) characteristic peaks  $\delta$  ppm; Ph: phenyl; Me: methyl; diMe: dimethyl; Et: ethyl; Pr: propyl; iPr: isopropyl; iBu: isobutyl; Pen: pentyl; cPr: cyclopropyl; Ac: acetyl; Cl: chloro; diCl: dichloro; CN: 15 cyano; F: fluoro; diF: difluoro; triF: trifluoro; NO<sub>2</sub>: nitro; MeO: methoxy; diMeO: dimethoxy; Br: bromo; diBr: dibromo; CF<sub>3</sub>: trifluoromethyl; AcO: acetoxy; MeOCO: methoxycarbonyl; Boc: tert-butoxycarbonyl; NH<sub>2</sub>: amino; PhCONH: benzoylamino; EtCONH: ethylcarbonylamino; Et<sub>2</sub>N: diethylamino; TBS: tert- 20 butyldimethylsilyl; biPh: biphenyl; Naph: naphthalene; Thiop: thiophene; Fu: furan; Py: pyridine; IM: imidazole; Pyrazi: pyrazine; Pipe: piperidine; Pyrazo: pyrazole; Pyrim: pyrimidine; Pyrr: pyrrole; Pyrroli: pyrrolidine; Mo: morpholine; Isoquin: isoquinoline; Isoind: isoindoline; Thiaz: 25 thiazole; Tr: triphenylmethyl; TEA: triethylamine; NMO: N-methylmorpholine oxide; TPAP tetrapropylammonium perruthenate;




Sa: addition salt; HCl: hydrochloride; Oxal: oxalate; MS4A: molecular sieves 4A.

Table 2




Compound No.	R <sup>2</sup>	R <sup>3</sup>	B	Compound No.	R <sup>2</sup>	R <sup>3</sup>	B
1a	H	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	Ph	7a	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	MeO	Ph
2a	H	MeOCOCH <sub>2</sub>	Ph	8a	H	Py-3-ylCH <sub>2</sub> NH	3-H <sub>2</sub> N-Ph
3a	H	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	Ph	9a	H	Py-3-ylCH <sub>2</sub> NH	Py-3-yl
4a	H	Me <sub>2</sub> NCOCH <sub>2</sub>	Ph	10a	H	4-O <sub>2</sub> N-PhCONH	3-H <sub>2</sub> N-Ph
5a	H	4-AcNH-PhCH <sub>2</sub> NH	Ph	11a	H	Me[MeO(CH <sub>2</sub> ) <sub>3</sub> ]NCH <sub>2</sub>	Ph
6a	H	MeCH(Ph)NH	Ph	12a	MeO	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	Ph

Table 3




Compound No.	R <sup>2</sup>	R <sup>3</sup>	A	B
13a	1,3-Thiaz-5-ylCH <sub>2</sub> NH	H	3,5-diF-Ph	3-Me-Ph
14a	O <sub>2</sub> N	Cl	3,5-diF-Ph	3-Me-Ph
15a	H	F	Ph	4-cPrNH-Ph
16a	MeO	MeO	3-HOOC-Ph	Ph
17a	H	H	3,5-diF <sub>3</sub> C-Ph	3-H <sub>2</sub> N-Ph
18a	4-F-PhCONH	H	Ph	Ph
19a	4-F-PhCONH	H	Ph	3,5-diF-Ph
20a	H	H	4-F-Ph	4-F-Ph

Table 4




Compound No.	A	B	Compound No.	A	B
21a	4-Cl-Ph	4-Cl-Ph	31a	3-MeO-Ph	3-MeO-Ph
22a	4-CN-Ph	4-CN-Ph	32a	3-Br-Ph	3-Br-Ph
23a	4-Me-Ph	4-Me-Ph	33a	3-Me-Ph	3-Me-Ph
24a	4-O <sub>2</sub> N-Ph	4-O <sub>2</sub> N-Ph	34a	3-Et-Ph	3-Et-Ph
25a	4-MeOCO-Ph	4-MeOCO-Ph	35a	Ph	3-F-Ph
26a	2-Cl-Ph	2-Cl-Ph	36a	3-H <sub>2</sub> N-Ph	3-H <sub>2</sub> N-Ph
27a	3-Cl-Ph	3-Cl-Ph	37a	3-(Py-3-ylCH <sub>2</sub> CONH)Ph	3,5-diF-Ph
28a	4-Cl-CH <sub>2</sub> -Ph	4-Cl-CH <sub>2</sub> -Ph	38a	4-(Mo-4-ylCH <sub>2</sub> )Ph	4-(Mo-4-ylCH <sub>2</sub> )Ph
29a	2-F-Ph	2-F-Ph	39a	3-OH-Ph	3-OH-Ph
30a	4-MeO-Ph	4-MeO-Ph	40a	3,5-diF-Ph	Py-3-ylCH <sub>2</sub> NHPh

Table 5




Compound No.	R <sup>2</sup>	Compound No.	R <sup>2</sup>
41a	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	81a	4-Cl-PhCH <sub>2</sub> NH
42a	MeOCOCH <sub>2</sub>	82a	2-Br-PhCH <sub>2</sub> NH
43a	Me[MeO(CH <sub>2</sub> ) <sub>3</sub> ]NCH <sub>2</sub>	83a	2-F-PhCH <sub>2</sub> NH
44a	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	84a	3-F-PhCH <sub>2</sub> NH
45a	Me <sub>2</sub> NCOCH <sub>2</sub>	85a	4-F-PhCH <sub>2</sub> NH
46a	4-AcNH-PhCH <sub>2</sub> NH	86a	2-HO-PhCH <sub>2</sub> NH
47a	MeCH(Ph)NH	87a	3-HO-PhCH <sub>2</sub> NH
48a	6-CF <sub>3</sub> -Py-3-ylCH <sub>2</sub> NH	88a	2-O <sub>2</sub> N-PhCH <sub>2</sub> NH
49a	4-tBuOCONH-Py-3-ylCH <sub>2</sub> NH	89a	3,5-diMeO-PhCH <sub>2</sub> NH
50a	2-Cl-Py-3-ylCH <sub>2</sub> NH	90a	2,5-diMeO-PhCH <sub>2</sub> NH
51a	4-H <sub>2</sub> N-Py-3-ylCH <sub>2</sub> NH	91a	2,3-diMeO-PhCH <sub>2</sub> NH
52a	6-Me-Py-2-ylCH <sub>2</sub> NH	92a	3,4-diF-PhCH <sub>2</sub> NH
53a	3-Cl-4-F <sub>3</sub> C-Py-2-ylCH <sub>2</sub> NH	93a	2,4-diF-PhCH <sub>2</sub> NH
54a	4,6-diMe-Py-2-ylCH <sub>2</sub> NH	94a	Fu-2-ylCH <sub>2</sub> NH
55a	5-CN-6-MeS-Py-2-ylCH <sub>2</sub> NH	95a	5-Me-Fu-2-ylCH <sub>2</sub> NH
56a	3,6-diCl-4-OH-Py-2-ylCH <sub>2</sub> NH	96a	4-iBu-PhCH <sub>2</sub> NH
57a	Py-2-ylCH <sub>2</sub> NH	97a	4-Br-PhCH <sub>2</sub> NH
58a	Py-4-ylCH <sub>2</sub> NH	98a	3-MeO-CO-PhCH <sub>2</sub> NH
59a	2,6-diCl-Py-4-ylCH <sub>2</sub> NH	99a	4-CN-PhCH <sub>2</sub> NH
60a	3,5-diOH-2-Me-Py-4-ylCH <sub>2</sub> NH	100a	3-PhCH <sub>2</sub> O-PhCH <sub>2</sub> NH
61a	Py-4-yl-CONH	101a	2-Cl-4-F-PhCH <sub>2</sub> NH
62a	3-MeO-CO-PhCONH	102a	2-Cl-5-HO-PhCH <sub>2</sub> NH
63a	4-(iPrNHCO)PhCH <sub>2</sub> NH	103a	3-Cl-4-MeO-PhCH <sub>2</sub> NH
64a	1-Me-IM-4-ylCH <sub>2</sub> NH	104a	3-Cl-6-O <sub>2</sub> N-PhCH <sub>2</sub> NH
65a	Py-2-ylCH <sub>2</sub> NH	105a	4-Cl-5-O <sub>2</sub> N-PhCH <sub>2</sub> NH
66a	6-Br-imidazo[1,2-a]Py-3-ylCH <sub>2</sub> NH	106a	2,3-diHO-PhCH <sub>2</sub> NH
67a	3-Cl-PhCH <sub>2</sub> NH	107a	2,4-diHO-PhCH <sub>2</sub> NH
68a	3-Br-PhCH <sub>2</sub> NH	108a	4,5-diHO-PhCH <sub>2</sub> NH
69a	4-Cl-PhCH <sub>2</sub> NH	109a	3-HO-4-MeO-PhCH <sub>2</sub> NH
70a	Naph-1-yl-CH <sub>2</sub> NH	110a	3-HO-5-O <sub>2</sub> N-PhCH <sub>2</sub> NH
71a	2-Me-PhCH <sub>2</sub> NH	111a	3-HO-4-O <sub>2</sub> N-PhCH <sub>2</sub> NH
72a	3-Me-PhCH <sub>2</sub> NH	112a	2-HO-6-MeO-PhCH <sub>2</sub> NH
73a	4-iPr-PhCH <sub>2</sub> NH	113a	4-MeO-PhCH <sub>2</sub> NH
74a	4-Et-PhCH <sub>2</sub> NH	114a	2-EtO-PhCH <sub>2</sub> NH
75a	2-MeO-PhCH <sub>2</sub> NH	115a	4-EtO-PhCH <sub>2</sub> NH
76a	4-MeO-Naph-1-yl-CH <sub>2</sub> NH	116a	4-MeO-Naph-1-yl-CH <sub>2</sub> NH
77a	4-MeO-3,6-diMe-PhCH <sub>2</sub> NH	185a	5-Me-IM-4-ylCH <sub>2</sub> NH
78a	3,5-diBr-6-HO-PhCH <sub>2</sub> NH	186a	IM-2-ylCH <sub>2</sub> NH
79a	2-CF <sub>3</sub> -PhCH <sub>2</sub> NH	187a	6-Me-Py-2-ylCH <sub>2</sub> NH
80a	3-Cl-PhCH <sub>2</sub> NH		

Table 6



Compound No.	R <sup>2</sup>	B	Compound No.	R <sup>2</sup>	B
117a	H	1H-IM-4-yl	151a	H	5-MeO-Py-3-yl
118a	H	Fu-2-yl	152a	H	2-H <sub>2</sub> N-Thiaz-4-yl
119a	H	3-PhNHCOPh	153a	H	1-(4-F-PhCH <sub>2</sub> )IM-4-yl
120a	H	3-H <sub>2</sub> N-5-F <sub>3</sub> C-Ph	154a	H	2-Me-Thiaz-4-yl
121a	O <sub>2</sub> N	3,5-diF-Ph	155a	H	5-Me-Py-3-yl
122a	O <sub>2</sub> N	2-Me-Ph	156a	Py-3-ylCH <sub>2</sub> NH	3-H <sub>2</sub> NPh
123a	O <sub>2</sub> N	3-F <sub>3</sub> C-O-Ph	157a	H	6-F <sub>3</sub> C-Py-3-yl
124a	O <sub>2</sub> N	3-Cl-Ph	158a	Py-3-ylCH <sub>2</sub> NH	Py-3-yl
125a	O <sub>2</sub> N	3,4-diMe-Ph	159a	H	1-Me-Pyrrol-3-yl
126a	O <sub>2</sub> N	4-MeO-Ph	160a	H	1,2,3-Thiadiazol-5-yl
127a	O <sub>2</sub> N	2-Cl-Ph	161a	4-NO <sub>2</sub> -PhCONH	3-H <sub>2</sub> N-Ph
128a	O <sub>2</sub> N	2,5-diF-Ph	162a	H	Pyrazine-2-yl
129a	O <sub>2</sub> N	2-F <sub>3</sub> C-Ph	163a	H	1-Me-benzoIM-5-yl
130a	O <sub>2</sub> N	3,5-diMe-Ph	164a	Py-3-ylCONH	3-Me-Ph
131a	O <sub>2</sub> N	2-F-Ph	165a	3-Cl-PhSO <sub>2</sub> NH	3-Me-Ph
132a	O <sub>2</sub> N	3,5-diMeO-Ph	166a	4-AcNH-PhCH <sub>2</sub> NH	3,5-diF-Ph
133a	O <sub>2</sub> N	5-Br-Py-3-yl	167a	4-AcNH-PhCH <sub>2</sub> NH	2-Me-Ph
134a	O <sub>2</sub> N	3-Br-Ph	168a	4-AcNH-PhCH <sub>2</sub> NH	4-F <sub>3</sub> C-O-Ph
135a	O <sub>2</sub> N	3-Me-Ph	169a	4-AcNH-PhCH <sub>2</sub> NH	3-F <sub>3</sub> C-O-Ph
136a	H <sub>2</sub> N	3-F-Ph	170a	4-AcNH-PhCH <sub>2</sub> NH	3-F <sub>3</sub> C-Ph
137a	H <sub>2</sub> N	4-Me-Ph	171a	4-AcNH-PhCH <sub>2</sub> NH	4-Cl-Ph
138a	H <sub>2</sub> N	4-F <sub>3</sub> C-O-Ph	172a	4-AcNH-PhCH <sub>2</sub> NH	3,4-diMe-Ph
139a	H <sub>2</sub> N	2-F <sub>3</sub> C-O-Ph	173a	4-AcNH-PhCH <sub>2</sub> NH	4-MeO-Ph
140a	H <sub>2</sub> N	4-Cl-Ph	174a	4-AcNH-PhCH <sub>2</sub> NH	3,5-diMeO-Ph
141a	H <sub>2</sub> N	3-Cl-Ph	175a	4-AcNH-PhCH <sub>2</sub> NH	4-F <sub>3</sub> C-Ph
142a	H <sub>2</sub> N	3,4-diMe-Ph	176a	3-Cl-PhCH <sub>2</sub> NH	4-F-Ph
143a	H <sub>2</sub> N	4-MeO-Ph	177a	4-HO-PhCH <sub>2</sub> NH	4-F-Ph
144a	H <sub>2</sub> N	2-Cl-Ph	178a	3-CN-PhCH <sub>2</sub> NH	2-MeO-Ph
145a	H <sub>2</sub> N	2-F <sub>3</sub> C-Ph	179a	3-Cl-PhCH <sub>2</sub> NH	2-MeO-Ph
146a	H <sub>2</sub> N	2-F-Ph	180a	4-HOOC-PhCH <sub>2</sub> NH	2-MeO-Ph
147a	H <sub>2</sub> N	3,5-diMeO-Ph	181a	4-HO-PhCH <sub>2</sub> NH	3-Me-Ph
148a	H <sub>2</sub> N	4-F <sub>3</sub> C-Ph	182a	2-Cl-PhCH <sub>2</sub> NH	3-Me-Ph
149a	H <sub>2</sub> N	3-Br-Ph	183a	3-Br-PhCH <sub>2</sub> NH	Ph
150a	H	5-Me-Py-3-yl	184a	4-Cl-PhCH <sub>2</sub> NH	Ph

Table 7



Compound No.	R <sup>2</sup>	B
188a	H	3-{Me[MeO(CH <sub>2</sub> ) <sub>2</sub> ]N}Ph
189a	H	2-H <sub>2</sub> N-CH <sub>2</sub> -Ph
190a	H	2-(1-HOOC-EtNH)Ph
191a	H	3-HOOC-Ph
192a	Pipe-1-yl	Ph
193a	H	4-H <sub>2</sub> NCO-IM-1-yl
194a	PhNHCO-diMe-C	Ph
195a	3-CN-PhNHCOCH <sub>2</sub>	Ph
196a	Py-4-ylCH <sub>2</sub> OCOCH <sub>2</sub>	Ph
197a	H	3-H <sub>2</sub> N-5-F-Ph
198a	3-F-PhCH <sub>2</sub> NHCH <sub>2</sub>	Ph
199a	F	4-cPr-NH-Ph
200a	Py-4-ylCONH	Ph
201a	3-MeOCOPhCH <sub>2</sub> CO	Ph
202a	6-F <sub>3</sub> C-Py-3-ylCH <sub>2</sub> NH	3,5-diF-Ph
203a	4-tBuOCONH-Py-3-ylCH <sub>2</sub> NH	3,5-diF-Ph
204a	2-Cl-Py-3-ylCH <sub>2</sub> NH	3,5-diF-Ph
205a	4-H <sub>2</sub> N-Py-3-ylCH <sub>2</sub> NH	3,5-diF-Ph
206a	6-Me-Py-2-ylCH <sub>2</sub> NH	3,5-diF-Ph
207a	3-Cl-4-F <sub>3</sub> C-Py-2-ylCH <sub>2</sub> NH	3,5-diF-Ph
208a	4,6-diMe-Py-2-ylCH <sub>2</sub> NH	3,5-diF-Ph
209a	5-CN-6-MeS-Py-2-ylCH <sub>2</sub> NH	3,5-diF-Ph
210a	3,6-diCl-4-OH-Py-2-ylCH <sub>2</sub> NH	3,5-diF-Ph
211a	Py-2-ylCH <sub>2</sub> NH	3,5-diF-Ph
212a	Py-4-ylCH <sub>2</sub> NH	3,5-diF-Ph
213a	2,6-diCl-Py-4-ylCH <sub>2</sub> NH	3,5-diF-Ph
214a	3,5-diOH-2-Me-Py-4-ylCH <sub>2</sub> NH	3,5-diF-Ph

Table 8

Compound No.	Str	Compound No.	Str	Compound No.	Str
215a		217a		219a	
216a		218a		222a	

The active ingredient of the invention and the compound of the invention or a pharmaceutically acceptable salt thereof can be employed solely as a pharmaceutical drug but usually, one or two or more of the active ingredients can be formulated by a generally used method using drug carriers, fillers and the like generally used in the art. Its administration may be either oral administration by tablets, pills, capsules, granules, powders, solutions and the like, or parenteral administration by intra-articular, intravenous, intramuscular, and the like injections, suppositories, eye drops, ophthalmic ointments, percutaneous solutions, ointments, percutaneous adhesive preparations, transmucosal solutions, transmucosal adhesive preparations, inhalations and the like.

The solid composition for use in the oral administration according to the present invention is used in the form of tablets, powders, granules and the like. In such a solid composition, one or more active ingredients are mixed with at least one inert diluent such as lactose, mannitol, glucose, hydroxypropylcellulose, microcrystalline cellulose, starch, polyvinyl pyrrolidone, magnesium aluminate metasilicate and/or the like. In accordance with the usual procedures, the composition may contain inert additives other than the diluent, for example, a lubricant such as magnesium stearate, a disintegrating agent such as calcium cellulose glycolate, a stabilizer such as lactose, and a solubilization assisting agent such as glutamic acid or aspartic acid. If

necessary, tablets or pills may be coated with a sugar or a film of a gastric or enteric coating substance, such as sucrose, gelatin, hydroxypropyl cellulose or hydroxypropylmethyl cellulose phthalate.

5           The liquid composition for oral administration includes pharmaceutically acceptable emulsions, solutions, suspensions, syrups, elixirs and the like and contains a generally used inert diluent, e.g., purified water or ethanol. In addition to the inert diluent, this composition may further  
10 contain an auxiliary agent such as a solubilizing agent, a moistening agent, a suspending agent or the like, as well as a sweetener, a flavor, an aromatic and a preservative.

          The injections for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions and  
15 emulsions. Examples of the aqueous solutions or suspensions include distilled water for injection and saline. Examples of the non-aqueous solutions or suspensions include propylene glycol, polyethylene glycol, vegetable oils such as olive oil, alcohols such as ethanol, polysorbate 80 (trade name) and the  
20 like. Such a composition may further contain auxiliary agents such as a tonicity agent, a preservative, a moistening agent, an emulsifier, a dispersing agent, a stabilizer (e.g., lactose) and a solubilization assisting agent (e.g., glutamic acid or aspartic acid). These compositions are sterilized,  
25 e.g., by filtration through a bacteria-retaining filter, blending of a germicide or irradiation. Alternatively, they may be used by firstly making into sterile solid compositions

and then dissolving or suspending them in sterile water or a sterile solvent for injection use prior to their use.

The transmucosal preparations such as transnasal preparations are used in the form of solid, liquid or semi-solid, and can be produced in accordance with hitherto known methods. For example, known pH adjusting agent, preservative, thickener and filler are optionally added and the preparations are formed into solid, liquid or semi-solid. The transnasal preparations are administered by means of a usual sprayer, nasal container, tube, intranasal insert or the like.

In the case of oral administration, suitable daily dose is usually about 0.001 to 100 mg/kg, preferably 0.1 to 30 mg/kg, more preferably 0.1 to 10 mg/kg-body weight, and the dose is divided into 1 or 2 to 4 doses per day. In the case of intravenous administration, suitable daily dose is about 0.0001 to 10 mg/kg-body weight, and the dose is divided into 1 to several doses per day. And, in the case of transmucosal preparations, about 0.001 to 100 mg/kg-body weight is divided into 1 to several doses per day. The dose may be appropriately determined for each case, depending on conditions, age, sex and the like.

**(Examples)**

The following will explain the invention further in detail based on Examples. The compounds of the invention are not limited to the compounds described in the following



Examples. In this connection, production methods of starting compounds are shown in Reference Examples.

#### Reference Example 1

To a boiling suspension of sodium hydride (60% in oil) (360 mg) in anhydrous THF (10 ml) was added dropwise a solution of acetophenone (720 mg) and ethyl 2-methylthiazol-4-carboxylate (1.20 g) in anhydrous THF (10 ml), followed by 10 minutes of heating under refluxing. After the reaction solution was cooled, a mixed solution of acetic acid (1 ml) and water (30 ml) was added thereto, and the resulting mixture was extracted with ethyl acetate. The extracted solution was washed with water and then dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: hexane:ethyl acetate = 4:1 (v/v)) to obtain 1-(2-methylthiazol-4-yl)-3-phenylpropan-1,3-dione (1.3 g, 88%) as yellow crystals. Hereinafter, the compounds of Reference Examples 2 to 10 were obtained similarly.

#### Reference Example 11

To a solution of the compound (674 mg) obtained in Reference Example 1 in DMF (8 ml) was added potassium carbonate (1.14 g), followed by 1 hour of stirring at room temperature. After carbon disulfide (283 mg) was added to the reaction solution and the resulting mixture was stirred at room temperature for 2 hours, methyl iodide (0.369 ml) was further added, followed by 1.5 hours of stirring at room temperature. Water was added to the reaction solution and the

resulting mixture was extracted with ethyl acetate. The extract solution was washed with water and then dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent (v/v): hexane:ethyl acetate = 4:1) to obtain 2-(bismethylsulfanylmethylene)-1-(2-methylthiazol-4-yl)-3-phenylpropan-1,3-dione (555 mg, 64%) as a yellow oily substance. Hereinafter, the compounds of Reference Examples 12 to 23 were obtained similarly.

10 Reference Example 24

A catalytic amount of ammonium chloride was added into 100 ml acetic acid solution containing terephthalaldicarboxaldehyde (1.34 g) and 3-methylrhodanine (1.53 g), followed by about 12 hours of heating at 110°C.

15 After cooling upon standing, the formed yellow crystals were collected by filtration, washed with an appropriate amount of ethanol-water (10:1), and dried to obtain 4-(3-methyl-4-oxo-2-thioxothiazolidin-5-ylidenmethyl)benzaldehyde (1.91 g).

Reference Example 25

20 Into a saturated ammonia-methanol solution (60 ml) was added 4-chloro-3-nitro-N-(pyridin-3-ylmethyl)benzenesulfonamide (2.77 g), followed by about 2 days of heating at 100°C in a sealed tube. After the reaction, the solvent was evaporated under reduced pressure and the formed  
25 yellow crystals were collected by filtration and dried to obtain 4-amino-3-nitro-N-(pyridin-3-

ylmethyl)benzenesulfonamide (2.51 g). Hereinafter, the compound of Reference Examples 26 was obtained similarly.

#### Reference Example 27

A catalytic amount of Raney nickel was added to an  
5 ethyl acetate-ethanol (1:1) solution (200 ml) containing the compound (2.49 g) obtained in Reference Example 25, followed by a reaction in the presence of hydrogen gas at ordinary temperature under ordinary pressure. After the reaction, a filtrate obtainable by removing the catalyst by filtration was  
10 evaporated under reduced pressure to obtain 3,4-diamino-N-(pyridin-3-ylmethyl)benzenesulfonamide (2.22 g).

#### Reference Example 28

Into a DMF solution (150 ml) containing 4-amino-3-nitrophenol (4.72 g) were added successively potassium  
15 carbonate (12.8 g), tetrabutylammonium iodide (0.56 g) and 3-chloromethylpyridine hydrochloride (5.42 g), followed by about 1.5 hours of heating at 60°C. The reaction solution was concentrated under reduced pressure, ice-water (250 ml) and 1M hydrochloric acid aqueous solution (30 ml) were poured  
20 thereto, and the formed crystals were collected by filtration and dried to obtain 2-nitro-4-(pyridin-3-ylmethoxy)aniline (7.39 g). Hereinafter, the compounds of Reference Examples 35 and 60 were obtained similarly.

#### Reference Example 29

25 Into an ethyl acetate-ethanol (1:1) solution (300 ml) containing the compound (3.68 g) obtained in Reference Example 28 was added 10% palladium on carbon, followed by a reaction

in the presence of hydrogen gas at ordinary temperature under ordinary pressure. After the reaction, a filtrate obtainable by removing the catalyst by filtration was evaporated under reduced pressure to obtain 4-(pyridin-3-ylmethoxy)benzene-1,2-diamine (3.23 g). Hereinafter, the compound of Reference Example 30 was obtained similarly.

#### Reference Example 31

Triphenylpyridin-3-ylmethylphosphonium chloride (1.95 g) was added into an ice-cooled DMF suspension (40 ml) containing sodium hydride (60% in oil) (0.26 g), followed by 30 minutes of stirring at room temperature. The reaction solution was cooled to 0°C and 4-chloro-3-nitrobenzaldehyde (1.04 g) was added portionwise thereto, followed by 2 hours of stirring at room temperature. To the residue formed by evaporating the reaction solution under reduced pressure was poured an appropriate amount of purified water, followed by extraction with ethyl acetate. Thereafter, the organic layer was dried over anhydrous magnesium sulfate. The crude product obtained by solvent evaporation was purified by silica gel column chromatography to obtain 3-[2-(4-chloro-3-nitrophenyl)vinyl]pyridine (0.86 g) from the fractions eluted with ethyl acetate-hexane (2:1 (v/v)).

#### Reference Example 32

To a DMF solution (10 mL) of (2-methyl-1H-benzimidazol-5-yl)methanol (811 mg) were added tert-butyldimethylsilyl chloride (904 mg) and imidazole (680 mg), followed by 2 hours of stirring at room temperature. The

reaction solution was concentrated, water was added thereto, the mixture was extracted with chloroform, and the extract was dried over anhydrous magnesium sulfate. The organic layer was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 10:1 (v/v)) to obtain 5-(tert-butyltrimethylsilyloxymethyl)-2-methyl-1H-benzimidazole (1305 mg, 94%).

#### Reference Example 33

Into an ethanol solution (100 ml) containing the compound (2.2 g) obtained in Reference Example 27 was added triethyl orthoacetate (3.21 g), followed by about 12 hours of heating under refluxing. Concentrated hydrochloric acid (1 ml) was added dropwise and the whole was heated under refluxing for another 2 hours. Then, the reaction solution was evaporated under reduced pressure. The residue was washed with cooled water (100 ml) containing saturated sodium hydrogen carbonate aqueous solution (10 ml), collected by filtration and dried to obtain 2-methyl-1H-benzimidazol-5-sulfonic acid (pyridin-3-ylmethyl)amide (1.94 g). Hereinafter, the compounds of Reference Examples 34 and 36 were obtained similarly.

#### Reference Example 37

(1) Benzoyl chloride (32.5 ml) was added dropwise to a mixture of 2-methyl-5-nitrobenzimidazole (12.5 g) and TEA (38.8 ml) in Diglyme (63 ml) at room temperature. The reaction mixture was stirred at 100°C for 1 hour. Water was

added to the reaction mixture cooled to room temperature and the whole was stirred for 45 minutes. The reaction mixture was extracted with chloroform, the organic layer was washed with water and dried over anhydrous sodium sulfate, and then the solvent was evaporated under reduced pressure. The obtained crude crystals were recrystallized from chloroform-n-hexane to obtain 2-(1-benzoyl-1H-5-nitrobenzimidazol-2-yl)-1-phenylvinyl benzoate (29.7 g, 86%).

(2) The compound obtained in (1) (29.7 g) and morpholine (15.8 g) were dissolved in methanol (90 ml), followed by 30 minutes of heating under refluxing. After the reaction mixture was cooled to room temperature, water was added thereto and the whole was stirred for 2 hours. The formed precipitate was collected by filtration, washed with cold water, and then dried to obtain 2-(1,3-dihydro-2H-5-nitrobenzimidazol-2-ylidene)-1-phenylethan-1-one (16.7 g, 84%). Hereinafter, the compounds of Reference Examples 38 to 54 and Reference Examples 61 to 64 were obtained similarly.

Reference Example 55

Into a 1,4-dioxane solution (250 ml) containing 2-amino-4-chlorothiazol-5-carbaldehyde (10.83 g) was added 4-(dimethylamino)pyridine (1 g). Then, a 1,4-dioxane solution (100 ml) containing di-tert-butyl dicarbonate (29 g) was gradually added dropwise under heating at 60°C, and then the whole was continued to stir for about 30 minutes. After the reaction solution was cooled upon standing, the solvent was evaporated under reduced pressure and an appropriate amount of

5% potassium hydrogen sulfate aqueous solution was poured to the thus obtained residue, followed by extraction with ethyl acetate. After the organic layer was washed with water and dried over anhydrous magnesium sulfate, the crude product  
5 formed by solvent evaporation was purified by silica gel column chromatography to obtain tert-butyl (4-chloro-5-formylthiazol-2-yl)-carbamate (10.73 g) as pale brown crystals from the fractions eluted with ethyl acetate-toluene (2:3 (v/v)).

10 Reference Example 56

Under an argon stream, a dioxane solution (10 ml) of tris-t-butylphosphine (240 mg) was added to a mixture of p-methoxyphenyl boric acid (4.364 g),  
tris(dibenzylideneacetone)dipalladium (452 mg), cesium  
15 carbonate (10.561 g), 5-chloro-2-nitroaniline (4505 mg) and dioxane (50 ml), followed by 2 hours and 10 minutes of heating at 85°C. After cooling to room temperature on standing, diethyl ether (500 ml) and chloroform (500 ml) were added thereto. After insoluble matter was removed by filtration,  
20 the filtrate was concentrated to obtain an aimed compound, 5-(4'-methoxyphenyl)-2-nitroaniline (6.5 g).

Reference Example 57

A catalytic amount of concentrated sulfuric acid was added dropwise to an acetic anhydride (55 ml) suspension of 5-  
25 (4'-methoxyphenyl)-2-nitroaniline (2.02 g), followed by 3 hours and 20 minutes stirring at 40°C. After cooling to room temperature, diethyl ether (200 ml) was added thereto and the

precipitated powder was collected by filtration to obtain N-(4'-methoxy-4-nitrobiphenyl-3-yl)acetamide (596 mg).

#### Reference Example 58

A mixture of N-(4'-methoxy-4-nitrobiphenyl-3-yl)acetamide (500 mg), acetic acid (6 ml) and iron powder (308 mg) was stirred at 100°C for 50 minutes, and then cooled to room temperature, and insoluble matter was removed by filtration using celite. A saturated sodium carbonate aqueous solution was added to the filtrate to render the liquid about pH 7, followed by extraction with chloroform. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure to obtain 5-(4'-methoxyphenyl)-2-methylbenzimidazole (320 mg).

#### Reference Example 59

To a dimethylformamide (20 ml) solution of 2-methylbenzimidazol-5-carboxylic acid (1.00 g) were added hydroxybenzotriazole (844 mg), 1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (1.21 g) and 4-methoxyphenylmethylaniline (1.33 g) at room temperature, and the reaction solution was stirred at room temperature for 18 hours. The reaction solution was concentrated under reduced pressure and the obtained residue was diluted with chloroform (20 ml). The organic layer was washed with saturated sodium hydrogen carbonate aqueous solution, water and saturated sodium chloride aqueous solution, and dried over anhydrous sodium sulfate. The residue obtained by solvent evaporation under reduced pressure was subjected to silica gel column



chromatography and eluted with chloroform-methanol (30:1 (v/v)) to obtain (4-methoxyphenylmethyl)amide of 2-methyl-1H-benzimidazol-5-carboxylic acid (1.27 g, 99%).

#### Example 1

5           5-Chloro-2-methylbenzimidazole (833 mg) was dissolved in Diglyme (4 ml), and TEA (2.43 ml) was added thereto. Benzoyl chloride (2.0 ml) was further added thereto, followed by 15 minutes of stirring at about 100°C. Water (0.1 ml) was added dropwise to the reaction solution and the whole was  
10           heated under stirring at 175°C for 10 minutes. After the reaction solution was cooled with air, water (15 ml) was added and the mixture was stirred, then the supernatant was decanted. Methanol (5 ml) was added to the residue and the precipitated crystals were collected by filtration, washed  
15           with cold methanol, and dried to obtain 2-(5-chloro-1,3-dihydro-2H-benzimidazol-2-yliden)-1,3-diphenylpropan-1,3-dione (706 mg, 38%) as pale yellow powdery crystals. Hereinafter, the compounds of Examples 2 to 25, 119 and 126 were obtained similarly.

#### 20           Example 26

          In a similar manner to Reference Example 37, 2-(1-benzoyl-1H-benzimidazol-2-yl)-1-phenylvinyl benzoate (26.8 g, 86%) was obtained in Step (1) and 2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-1-phenylethan-1-one (11.9 g, 84%) in  
25           Step (2).

(3)       To a mixture of 3,5-difluorobenzoyl chloride (1.67 g) and pyridine (8.5 ml) was added portionwise the compound (1.01

g) obtained in the above (2), followed by 3 hours of stirring at room temperature. Water was added to the reaction mixture, followed by extraction with chloroform. The obtained organic layer was washed with water, saturated ammonium chloride aqueous solution and saturated brine, and dried over anhydrous sodium sulfate, and then the solvent was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain 2-[1-(3,5-difluorobenzoyl)-1H-benzimidazol-2-yl]-1-phenylvinyl 3,5-difluorobenzoate (1.45 g, 65%) as yellowish white powdery crystals.

(4) The compound (931 mg) obtained in (3) and 3,5-difluorobenzoic acid (570 mg) were dissolved in Diglyme (2.5 ml), followed by 20 minutes of stirring at 175°C. Water was added to the reaction mixture cooled to room temperature, the mixture was extracted with chloroform, the organic layer was washed with water and dried over anhydrous sodium sulfate, and then the solvent was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain yellow powdery crystals from the fractions eluted with chloroform-n-hexane. The crystals were recrystallized from methanol to obtain 1-(3,5-difluorophenyl)-2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-3-phenylpropan-1,3-dione (603 mg, 89%). Hereinafter, the compounds of Examples 27 to 39, 117, 118, 120 to 125, 127 to 166, 425, 431 and 446 were obtained similarly.

**Example 40**

A mixture of the compound (317 mg) obtained in Example 35, platinum (IV) oxide (30 mg) and ethyl acetate (30 ml) was stirred at room temperature for 23 hours under hydrogen atmosphere. After a black powder was removed by filtration, the filtrate was concentrated and the obtained residue was treated with 4M hydrogen chloride-ethyl acetate solution to obtain 1-(3-aminophenyl)-3-(3,5-difluorophenyl)-2-(1,3-dihydro-2H-benzimidazol-2-ylidene)propane-1,3-dione hydrochloride (245 mg, 76%) as a green powder. Hereinafter, the compounds of Examples 41 to 43, 167 to 203, 411, 412 and 432 were obtained similarly.

**Example 44**

The compound (200 mg) obtained in Example 43 was dissolved in pyridine (2 ml), and propionyl chloride (58 mg) was added dropwise thereto under ice cooling. The reaction temperature was raised to room temperature and the whole was stirred for 1 hour. Water was added to the reaction mixture, followed by extraction with chloroform. The obtained organic layer was washed with water and saturated brine and dried over anhydrous sodium sulfate, and then the solvent was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain 3'-[2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-3-oxo-3-phenylpropanoyl]propananilide (204 mg, 88%) as a yellow foamed powder from the fractions eluted with chloroform-methanol. Hereinafter, the compounds of Examples 45 to 78, 204 to 237,

416 to 420, 430, 433, 440 to 442 and 449 were obtained similarly.

#### Example 79

The compound (162 mg) obtained in Example 39 was dissolved in DMF (10 ml), and 4-(2-aminoethyl)pyridine (348 mg), potassium carbonate (591 mg) and potassium iodide (473 mg) were added thereto, followed by 7 hours stirring at room temperature. Ethyl acetate and water were added to the reaction mixture and the organic layer was separated. The obtained organic layer was washed with water and saturated brine and dried over anhydrous sodium sulfate, and then the solution was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography and the fractions eluted with chloroform were dissolved into ethyl acetate, then ethanolic hydrochloric acid was added thereto. The formed crystals were filtered to obtain 2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-1-phenyl-3-(3-((2-pyridin-4-ylethyl)amino)methyl)phenyl)propane-1,3-dione hydrochloride (417 mg, 51%) as a pale pink powder. Hereinafter, the compounds of Examples 80, 81 and 450 were obtained similarly.

#### Example 82

The compound (343 mg) obtained in Example 30 was dissolved in ethanol (8 ml), and morpholine (0.4 ml) was added thereto, followed by 2 hours of heating under refluxing. The reaction solution was cooled and then evaporated under reduced pressure. Chloroform and water were added to the obtained residue and the organic layer was separated. The obtained

organic layer was washed with water and saturated brine and dried over anhydrous sodium sulfate, and then the solution was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain 2-  
5 (1,3-dihydro-2H-benzimidazol-2-ylidene)-1-(3-hydroxyphenyl)-3-phenylpropane-1,3-dione (125 mg, 41%) as a yellow powder from the fractions eluted with chloroform.

#### Example 83

The compound (450 mg) obtained in Example 40 was  
10 dissolved in benzene (30 ml), and 4-formylimidazole (121 mg) and a catalytic amount of p-toluenesulfonic acid were added thereto, followed by stirring at room temperature for 3 hours, at 50°C for 2.5 hours and under heating with refluxing for 3.5 hours. The residue after solvent evaporation was dissolved in  
15 methanol (25 ml), and sodium borohydride (44 mg) was added under ice cooling, followed by 1 hour and 40 minutes of stirring. Water, chloroform and isopropanol were added to the reaction solution and the organic layer was separated. The residue obtained by concentrating the obtained organic layer  
20 was subjected to silica gel column chromatography to obtain 1-(3,5-difluorophenyl)-2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-3-{3-[(1H-imidazol-4-ylmethyl)amino]phenyl}propane-1,3-dione from the fractions eluted with chloroform-methanol. This compound was converted into a hydrochloride salt using 4M  
25 hydrogen chloride-ethyl acetate solution to obtain 1-(3,5-difluorophenyl)-2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-3-{3-[(1H-imidazol-4-ylmethyl)amino]phenyl}propane-1,3-dione

hydrochloride (159 mg, 27%) as a pale blue powder.

Hereinafter, the compounds of Examples 395 to 396 were obtained similarly.

#### Example 84

5 To a methylene chloride (3 mL) solution of the compound (180 mg) obtained in Example 43 was added pyridin-3-aldehyde (60 mg) and acetic acid (153 mg), and sodium triacetoxymethylborohydride (215 mg) was further added thereto under ice cooling, followed by 15 hours of stirring at room  
10 temperature. A saturated sodium hydrogen carbonate aqueous solution was added to the reaction mixture and the mixture was extracted with methylene chloride. After washing with water and saturated brine, the extract was dried over anhydrous magnesium sulfate. The organic layer was concentrated under  
15 reduced pressure and the obtained residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 30:1 (v/v)). The purified product was dissolved into chloroform (3 mL) and subjected to salt formation with 4M-HCl-ethyl acetate solution to obtain 2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-1-phenyl-3-(3-(pyridin-3-  
20 ylmethylamino)phenyl)propane-1,3-dione hydrochloride (186 mg, 76%). Hereinafter, the compounds of Examples 85 to 100, 238 to 393, 410, 413 to 415, 421 to 424, 426, 428, 429, 435 to 437, 439 and 443 to 445 were obtained similarly.

#### 25 Example 101

The compound (512 mg) obtained in Reference Example 11 was dissolved in ethanol (6 ml), and 1,2-phenylenediamine (237

mg) was added thereto, followed by 13 hours of heating under refluxing. The reaction solution was cooled and the formed crystals were collected by filtration and washed with methanol to obtain 2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-1-(2-methylthiazol-4-yl)-3-phenylpropane-1,3-dione (171 mg, 32%) as a yellow powder. Hereinafter, the compounds of Examples 102 to 111, 397 and 398 were obtained similarly.

#### Example 112

(1) Using the compound obtained in Reference Example 19, 1-(5-benzyloxypyridin-3-yl)-3-(3,5-difluorophenyl)-2-(1,3-dihydro-2H-benzimidazol-2-ylidene)propane-1,3-dione was obtained in a similar manner to Example 101.

(2) The compound (121 mg) obtained in (1) was dissolved in ethanol (6 ml), and 10% palladium on carbon (160 mg) was added thereto, followed by 21 hours of vigorous stirring under hydrogen atmosphere. The catalyst was removed by filtration, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 10:1 (v/v)). The eluate was recrystallized from chloroform-methanol-hexane to obtain 1-(3,5-difluorophenyl)-2-(1,3-dihydro-2H-benzimidazol-2-ylidene)-3-(5-hydroxypyridin-3-yl)propane-1,3-dione (61 mg, 62%) as yellow crystals.

#### Example 113

The compound (150 mg) obtained in Example 20 was dissolved in dichloromethane (4 ml) under argon atmosphere, and 1.0M boron tribromide-methylene chloride solution (1.25

ml) was added dropwise thereto under ice cooling. After 1 hour of stirring at 0°C, the reaction temperature was raised to room temperature and the whole was further stirred for another 4 hours. Methanol (0.5 ml) was added to the reaction mixture under ice cooling, followed by 40 minutes of stirring. Then, chloroform and water were added thereto, and the organic layer was separated. The resulting organic layer was washed with water and saturated brine and dried over anhydrous sodium sulfate, and then the solution was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain 2-(5-hydroxy-1,3-dihydro-2H-benzimidazol-2-ylidene)-1,3-bis(3-methylphenyl)propane-1,3-dione (39 mg, 27%) as an orange powder from the fractions eluted with chloroform.

#### Example 114

Benzoyl chloride (1.68 g) was added to a mixture of 1,2-dimethylbenzimidazole (0.5 g) and TEA (1.21 g) in Diglyme (5 ml) at room temperature. The reaction mixture was stirred at 120°C for 1 hour and successively at 150°C for 6 hours.

Water was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was washed with water and dried over anhydrous magnesium sulfate, and then the solvent was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain crude crystals from the fractions eluted with chloroform. The crystals were recrystallized from ethyl acetate to obtain 2-(1-methyl-1H-benzimidazol-2-yl)-1,3-



diphenylpropane-1,3-dione (0.81 g). Hereinafter, the compound of Example 115 was obtained similarly.

#### Example 116

- (1) The compound (1.01 g) obtained in Example 3 and N-bromosuccinimide (609 mg) were dissolved in tetrachloromethane (14 ml), and azobisisobutyronitrile (47 mg) was added thereto, followed by 1 hour of heating under refluxing. After the reaction solution was cooled, the precipitated crystals were collected and dried to obtain 2-(1-bromo-5-methyl-1,3-dihydro-2H-benzimidazol-2-ylidene)-1,3-diphenylpropane-1,3-dione (1.22 g, 99%) as cream-colored powdery crystals.
- (2) The compound obtained in (1) (400 mg), potassium carbonate (153 mg) and diethylamine (0.115 ml) were dissolved in DMF (4 ml), followed by 4.5 hours of stirring at room temperature.
- The reaction mixture was poured into water and extracted with chloroform. The resulting organic layer was washed with water and saturated brine and dried over anhydrous sodium sulfate, and then the solution was evaporated under reduced pressure. The obtained residue was subjected to silica gel column chromatography to obtain a yellow oily substance from the fractions eluted with chloroform. To a solution of this substance dissolved in chloroform (1 ml) was added dropwise 4M-hydrogen chloride-ethyl acetate solution under ice cooling, followed by 30 minutes of stirring at room temperature. The precipitated crystals were collected by filtration, washed with chloroform, and dried to obtain 2-(1diethylamino-5-methyl-1,3-dihydro-2H-benzimidazol-2-ylidene)-1,3-

diphenylpropane-1,3-dione hydrochloride (203 mg, 48%) as a pale yellow powder.

#### Example 394

To a THF solution (5 ml) of 3-{2-[5-(4-acetylaminobenzylamino)-1,3-dihydro-2H-benzimidazol-2-ylidene]-3-(3,5-difluorophenyl)-3-oxopropionyl}phenyl acetate (123 mg) was added 1M sodium hydroxide aqueous solution (0.5 ml), followed by 24 hours of stirring. A saturated ammonium chloride aqueous solution was added thereto, followed by extraction with ethyl acetate and drying over anhydrous magnesium sulfate. The organic layer was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 20:1 (v/v)) to obtain N-[4-((2-[1-(3,5-difluorobenzoyl)-2-(3-hydroxyphenyl)-2-oxoethylidene]-2,3-dihydro-1H-benzimidazol-5-ylamino)methyl)phenyl]acetamide (98 mg, 86%). Hereinafter, the compound of Example 438 was obtained similarly.

#### Example 399

An ethanol solution (10 ml) containing the compound (0.23 g) obtained in Example 127 was cooled to -15°C and 90% sodium borohydride (30 mg) was added thereto, followed by 1 hour of stirring at the same temperature. Appropriate amounts of purified water and saturated brine were poured into the reaction solution, followed by extraction with ethyl acetate. The organic layer was dried and concentrated and the obtained residue was purified by silica gel column chromatography to

obtain 1-(3,5-difluorophenyl)-2-[5-(1-hydroxyethyl)-1,3-dihydro-2H-benzimidazol-2-ylidene]-3-phenylpropane-1,3-dione (90 mg) from the fractions eluted with chloroform-methanol (50:1 (v/v)). Hereinafter, the compound of Example 400 was  
5 obtained similarly.

#### Example 401

Into an acetic acid solution (40 ml) containing the compound (0.77 g) obtained in Example 132 was added 10% palladium on carbon (80 mg), followed by stirring under  
10 hydrogen gas atmosphere at ordinary temperature under ordinary pressure. After removal of the catalyst by filtration, the solvent was evaporated under reduced pressure, followed by extraction with ethyl acetate. The ethyl acetate layer was washed with appropriate amounts of sodium hydrogen carbonate  
15 aqueous solution and saturated brine successively and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to obtain 1-(3,5-difluorophenyl)-2-(5-hydroxy-1,3-dihydro-2H-benzimidazol-2-ylidene)-3-phenylpropane-1,3-dione (0.58 g).

#### 20 Example 402

Into an acetonitrile solution (4 ml) containing the compound (100 mg) obtained in Example 401 and (3-chloromethyl)pyridine hydrochloride (50 mg) were added potassium carbonate (83 mg) and a catalytic amount of sodium  
25 iodide successively, followed by 3.5 hours of heating at 80°C. After solvent evaporation, an appropriate amount of purified water was poured, followed by extraction with ethyl acetate.

After drying over anhydrous magnesium sulfate and concentration, the resulting residue was purified by silica gel column chromatography to obtain yellow foam (43 mg) from the fractions eluted with chloroform-methanol (200:1 (v/v)).

5 This substance was dissolved into acetone (2 ml), and oxalic acid (16 mg) was added thereto, followed by stirring. The resulting crystals were collected by filtration to obtain 1-(3,5-difluorophenyl)-2-[5-(pyridin-3-ylmethoxy)-1,3-dihydro-2H-benzimidazol-2-ylidene]-3-phenylpropane-1,3-dione oxalate  
10 (35 mg).

#### Example 403

Into a dichloromethane solution (5 ml) containing the compound (0.13 g) obtained in Example 136 was added 80% mcpba (0.14 g), followed by 2 hours of stirring at room temperature.

15 The reaction solution was washed with sodium hydrogen sulfite aqueous solution and sodium hydrogen carbonate aqueous solution successively, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the resulting residue was recrystallized  
20 from ethyl acetate-hexane (1:1 (v/v)) to obtain 1-(3,5-difluorophenyl)-3-phenyl-2-(5-phenylmethanesulfonyl-1,3-dihydro-2H-benzimidazol-2-ylidene)propane-1,3-dione (94 mg).

#### Example 404

A chloroform solution (3 ml) containing the compound  
25 (145 mg) obtained in Example 239 was cooled with ice and 80% mcpba (80 mg) was added thereto, followed by 1 hour of stirring at room temperature. The reaction solution was

washed with sodium hydrogen sulfite aqueous solution and sodium hydrogen carbonate aqueous solution successively, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. Then, the resulting  
5 residue was purified by silica gel column chromatography to obtain 1-(3,5-difluorophenyl)-3-phenyl-2-{5-[(1-oxidopyridin-3-ylmethyl)amino]-1,3-dihydro-2H-benzimidazol-2-ylidene}propane-1,3-dione (92 mg) from the fractions eluted with chloroform-methanol (100:1 (v/v)).

10 **Example 405**

Into a THF/water = 1:1 solution (4 ml) of the compound (62 mg) obtained in Example 131 was added acetic acid (2 ml), followed by 4 hours of stirring at room temperature. Thereto was added a saturated sodium hydrogen carbonate aqueous  
15 solution, followed by extraction with chloroform and drying over anhydrous magnesium sulfate. The organic layer was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 20:1 (v/v)) to obtain 1-(3,5-  
20 difluorophenyl)-2-(5-hydroxymethyl-1,3-dihydro-2H-benzimidazol-2-ylidene)-3-phenylpropane-1,3-dione (43 mg, 89%).

**Example 406**

Into an ethanol solution (6 ml) containing the  
25 compound (0.30 g) obtained in Example 393 was added platinum oxide (0.03 g), followed by 8.5 hours of stirring at room temperature under hydrogen atmosphere. After solid matter in

the reaction solution was removed by filtration and the filtrate was evaporated under reduced pressure, the resulting residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 100:3 (v/v)) to obtain 2-{5-  
5 [(1H-benzimidazol-5-ylmethyl)amino]-1,3-dihydrobenzimidazol-2-ylidene}-1-(3,5-difluorophenyl)-3-phenylpropane-1,3-dione (24 mg, 12%).

#### Example 407

Into an acetonitrile solution (5 ml) containing the  
10 compound (160 mg) obtained in Example 196 was added phenyl isothiocyanate (60 mg), followed by 5 hours of stirring at room temperature. The formed crystals were collected by filtration, washed with a small amount of diethyl ether, and dried to obtain 1-{2-[1-(3,5-difluorobenzoyl)-2-oxo-2-  
15 phenylethylidene]-2,3-dihydro-1H-benzimidazol-5-yl}-3-phenylthiourea (0.19 g).

#### Example 408

To an acetonitrile solution (10 ml) of nicotinoyl chloride hydrochloride (356 mg) were added sodium azide (325  
20 mg) and triethylamine (0.836 ml), followed by 1.5 hours of stirring under ice cooling. Thereto was added water, followed by extraction with diethyl ether. After drying with anhydrous magnesium sulfate, the organic layer was concentrated under reduced pressure and toluene (10 ml) was added to the obtained  
25 residue, followed by 1 hour of heating under refluxing. After cooling to room temperature, an acetonitrile solution (5 ml) of the compound (255 mg) obtained in Example 196 was added and

the whole was stirred for 18 hours at room temperature. The reaction solution was concentrated under reduced pressure and the obtained residue was purified by silica gel column chromatography (eluent: chloroform:methanol = 10:1 (v/v)) and recrystallized (chloroform:methanol:hexane) to obtain 1-{2-[1-(3,5-difluorobenzoyl)-2-oxo-2-phenylethylidene]-2,3-dihydro-1H-benzimidazol-5-yl}-3-pyridin-3-ylurea (108 mg, 42%).

#### Example 409

Into a dichloromethane solution (5 ml) of the compound (174 mg) obtained in Example 405 were added NMO (100 mg) and MS4A, followed by 10 minutes of stirring at room temperature. TPAP (8 mg) was further added thereto, followed by 30 minutes of stirring at room temperature. The reaction solution was purified by silica gel column chromatography (eluent: chloroform:methanol = 40:1 (v/v)) to obtain 2-[1-(3,5-difluorobenzoyl)-2-oxo-2-phenylethylidene]-2,3-dihydro-1H-benzimidazol-5-carbaldehyde (76 mg, 44%).

#### Example 427

To a solution of N-{2-[1-benzoyl-2-(3,5-difluorophenyl)-2-oxoethylidene]-2,3-dihydro-1H-benzimidazol-5-yl}-4-methylbenzosulfonamide (376 mg), dichloromethane (40 ml) and methanol (10 ml) was added 2M trimethylsilyldiazomethane hexane solution (1.0 ml), and the reaction solution was stirred at room temperature for 15 hours. The solvent was evaporated under reduced pressure and the obtained residue was subjected to silica gel column chromatography and eluted with chloroform to obtain N-{2-[1-

benzoyl-2-(3,5-difluorophenyl)-2-oxoethylidene]-2,3-dihydro-1H-benzimidazol-5-yl}-N,4-dimethylbenzosulfonamide (310 mg). The obtained crude crystals were recrystallized from ethyl acetate-hexane to obtain crystals (181 mg, 47%).

5   **Example 434**

To a solution of 2-(5-amino-1,3-dihydro-2H-benzimidazol-2-ylidene)-1-(3,5-difluorophenyl)-3-phenylpropane-1,3-dione (400 mg) and ethanol (10 ml) was added hydroxymethylbenzotriazole (168 mg) at room temperature, and  
10   the reaction solution was stirred at room temperature for 20 hours. The reaction solution was filtered and the resulting solid matter was dissolved in THF (10 ml). Thereto was added sodium borohydride (78 mg) at room temperature and the reaction mixture was stirred at room temperature for 3 hours.  
15   The reaction solution was diluted with ethyl acetate (10 ml) and the organic layer was washed with saturated sodium hydrogen carbonate aqueous solution, water and saturated sodium chloride aqueous solution. The organic layer was dried over anhydrous sodium sulfate and the solvent was evaporated  
20   under reduced pressure. The obtained residue was subjected to silica gel column chromatography and eluted with chloroform-methanol (100:1 (v/v)) to obtain 1-(3,5-difluorophenyl)-2-[5-methylamino-1,3-dihydro-2H-benzimidazol-2-ylidene]-3-phenylpropane-1,3-dione (163 mg, 48%).

25   **Example 447**

Into a dichloromethane solution containing 2-[5-(4-aminobenzylamino)-1,3-dihydro-2H-benzimidazol-2-ylidene]-1-



(3,5-difluorophenyl)-3-(3-methylphenyl)propane-1,3-dione (0.28 g) and 2-chloro-1-methylpyridinium iodide (0.17 g) were added N,N-diisopropylethylamine (0.23 ml) and N,N'-(di-tert-butoxycarbonyl)thiourea (0.18 g) successively, followed by about 2 days of stirring at room temperature. The reaction solution was washed with an appropriate amount of purified water and then the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated and the obtained residue was subjected to silica gel column chromatography to obtain N,N'-(di-tert-butoxycarbonyl)-N"-[4-((2-[1-(3,5-difluorobenzoyl)-2-oxo-2-(3-methylphenyl)eth-(z)-ylidene]-2,3-dihydro-1H-benzimidazol-5-ylamino)methyl)phenyl]guanidine (0.31 g) from the fractions eluted with ethyl acetate-hexane (1:2 (v/v)).

#### Example 448

Into an ethyl acetate solution (3 ml) containing the compound (0.3 g) obtained in Example 447 was added dropwise 4M hydrogen chloride-ethyl acetate solution (3 ml), followed by about 2.5 hours of stirring at room temperature. The resulting white crystals were collected by filtration to obtain N-[4-((2-[1-(3,5-difluorobenzoyl)-2-oxo-2-(3-methylphenyl)eth-(z)-ylidene]-2,3-dihydro-1H-benzimidazol-5-ylamino)methyl)phenyl]guanidine hydrochloride (0.21 g).


The following tables show compounds obtained in the above Reference Examples and Examples and physicochemical properties thereof.

Table 9



Rex	A	B	DAT
1	Ph	2-Me-1,3-Thiaz-4-yl	FA:246
2	Ph	3-Mo-1-yl(CH <sub>2</sub> ) <sub>2</sub> O	FA:354
3	Ph	3-Me <sub>2</sub> N-Ph	FA:268
4	Ph	Me(PhCH <sub>2</sub> )NCH <sub>2</sub>	FA:358
5	Py-3-yl	3,5-diF-Ph	FA:262
6	Ph	3,5-diF-Ph	FA:261
7	5-PhCH <sub>2</sub> O-Py-3-yl	3,5-diF-Ph	FA:368
8	5-Me-Py-3-yl	3,5-diF-Ph	FA:276
9	1-Me-benzoIM-5-yl	3,5-diF-Ph	FA:315
10	6-Me-Py-3-yl	3,5-diF-Ph	FA:276

Table 10




Rex	A	B	DAT (FA:)	Rex	A	B	DAT (FA:)
11	Ph	2-Me-1,3-Thiaz-4-yl	350	17	Py-3-yl	3,5-diF-Ph	366
12	Ph	Py-4-yl	330	18	Ph	3,5-diF-Ph	261
13	Ph	Py-3-yl	330	19	5-PhCH <sub>2</sub> O-Py-3-yl	3,5-diF-Ph	368
14	Ph	3-Mo-1-yl(CH <sub>2</sub> ) <sub>2</sub> O	458	20	5-Me-Py-3-yl	3,5-diF-Ph	380
15	Ph	3-Me <sub>2</sub> N-Ph	371	21	1-Me-benzoIM-5-yl	3,5-diF-Ph	419
16	Ph	Me(PhCH <sub>2</sub> )NCH <sub>2</sub>	462	22	6-Me-Py-3-yl	3,5-diF-Ph	380

Table 11


Re x	Str	DAT	Rex	Str	DAT
23		FA:251	55		FA:263
24		EI:263			

Table 12




Rex	R <sup>2</sup>	R <sup>a</sup>	R <sup>b</sup>	DAT	Rex	R <sup>2</sup>	R <sup>a</sup>	R <sup>b</sup>	DAT
25	Py-3-ylCH <sub>2</sub> NHSO <sub>2</sub>	NO <sub>2</sub>	NH <sub>2</sub>	FA:309	30	Py-3-yl(CH <sub>2</sub> ) <sub>2</sub>	NH <sub>2</sub>	NH <sub>2</sub>	FA:214
26	Py-3-ylCH=CH	NO <sub>2</sub>	NH <sub>2</sub>	FA:242	31	Py-3-ylCH=CH	NO <sub>2</sub>	Cl	FA:261
27	Py-3-ylCH <sub>2</sub> NHSO <sub>2</sub>	NH <sub>2</sub>	NH <sub>2</sub>	FA:279	56	4-MeO-Ph	NO <sub>2</sub>	NH <sub>2</sub>	FN:243
28	Py-3-ylCH <sub>2</sub> O	NO <sub>2</sub>	NH <sub>2</sub>	FA:246	57	4-MeO-Ph	NO <sub>2</sub>	NHAc	FN:285
29	Py-3-ylCH <sub>2</sub> O	NH <sub>2</sub>	NH <sub>2</sub>	FA:216					

Table 13




Rex	R²	DAT	Rex	R²	DAT
32	TBS-OCH₂	FA:277	36	Py-3-yl(CH₂)₂	FA:238
33	Py-3-ylCH₂NHSO₂	FA:303	58	4-MeO-Ph	FA:239
34	Py-3-ylCH₂O	FA:240	59	4-MeO-Ph(Me)NCO	FA:296
35	4-O₂N-PhCH₂O	FA:284	60	PhCH₂O-CO	FA:267

Table 14




Rex	R²	R³	A	DAT (FA:)	Rex	R²	R³	A	DAT (FA:)
37	O₂N	H	Ph	282	48	PhCH₂S	H	Ph	359
38	O₂N	H	4-MeO-Ph	312	49	HOCH₂	H	Ph	267
39	O₂N	H	2-MeO-Ph	312	50	TBS-OCH₂	H	Ph	381
40	O₂N	H	2-Cl-Ph	316	51	Py-3-ylCH₂NHSO₂	H	Ph	407
41	O₂N	H	2,3-diMeO-Ph	342	52	Py-3-ylCH₂O	H	3,5-diF-Ph	380
42	O₂N	H	Thiop-2-yl	286	53	4-O₂N-PhCH₂O	H	3,5-diF-Ph	424
43	O₂N	H	3,5-diF-Ph	318	54	Py-3-yl(CH₂)₂	H	3,5-diF-Ph	378
44	O₂N	Cl	3,5-diF-Ph	352	61	4-MeO-Ph	H	3-Me-Ph	357
45	H	H	3,5-diF-Ph	273	62	O₂N	H	3-Me-Ph	296
46	Ac	H	Ph	279	63	4-MeO-Ph(Me)NCO	H	3,5-diF-Ph	436
47	PhCH₂O	H	Ph	343	64	PhCH₂O-CO	H	3,5-diF-Ph	407

Table 15




EX	R <sup>2</sup>	R <sup>3</sup>	A	B	Sa	DAT
1	Cl	H	Ph	Ph	—	FA: 375
2	O <sub>2</sub> N	H	Ph	Ph	—	FA: 386
3	Me	H	Ph	Ph	—	FA: 355
4	H	H	3-F-Ph	3-F-Ph	—	FA:377,N1:7.30-7.33(2H,m), 7.74-7.76(2H,m),13.15 (2H,s)
5	H	H	3,4-diCl-Ph	3,4-diCl-Ph	—	FA: 479
6	H	H	Fu-2-yl	Fu-2-yl	—	FA: 321
7	H	H	Thiop-2-yl	Thiop-2-yl	—	FA: 353
8	H	H	2-MeO-Ph	2-MeO-Ph	—	FA: 401
9	H	H	3-O <sub>2</sub> N-Ph	3-O <sub>2</sub> N-Ph	—	FA:431,N1:7.80-7.82(2H,m), 8.00-8.01(2H,m), 13.28(2H,s)
10	Me	Me	Ph	Ph	—	FA: 369
11	H	H	3-F <sub>3</sub> C-Ph	3-F <sub>3</sub> C-Ph	—	FA: 477
12	H	H	3-MeOCO-Ph	3-MeOCO-Ph	—	FA: 457
13	H	H	3-Cl-CH <sub>2</sub> -Ph	3-Cl-CH <sub>2</sub> -Ph	—	FA:437,N1:4.59(4H,s),7.73-7.76 (2H,m), 13.13(2H,s)
14	F	H	Ph	Ph	—	FA: 359
15	H	H	3-CN-Ph	3-CN-Ph	—	FA: 391
16	H	H	3-(PhCO)Ph	3-(PhCO)Ph	—	FA: 549
17	H	H	3-AcO-Ph	3-AcO-Ph	—	FA: 457
18	H	H	4-iPr-Ph	4-iPr-Ph	—	N1:1.03(d,6H,J=9),2.68(m,1H), 13.11(m,2H)
19	F	H	3-Me-Ph	3-Me-Ph	—	FA:387,N1:2.11(6H,s),6.17-7.18(9H,m),7.50-7.73(2H,m), 13.14-13.19(2H,m)
20	MeO	H	3-Me-Ph	3-Me-Ph	—	FA: 399
21	H	H	3,5-diF-Ph	3,5-diF-Ph	—	FA: 413
173	H <sub>2</sub> N	Cl	3,5-diF-Ph	3-Me-Ph	—	FA:440
429	4-AcNH-PhCH <sub>2</sub> NH	Cl	3,5-diF-Ph	3-Me-Ph	—	FA:587

Table 16




EX	R <sup>2</sup>	A	B	Sa	DAT
22	PhCOCH <sub>2</sub> OCO	Ph	Ph	—	FA: 503
23	PhCO	Ph	Ph	—	FA: 445
26	H	Ph	3,5-diF-Ph	—	FA:377,N1:7.30-7.34(4H,m),7.74-7.76(2H,m), 13.15(2H,s)
27	H	3-Me-Ph	Ph	—	FA:355,N1:2.10(3H,s),7.72-7.74(2H,m), 13.11(2H,s)
28	H	3-O <sub>2</sub> N-Ph	Ph	—	FA:386,N1:7.74-7.79(3H,m),7.93-7.98(2H,m), 13.20(2H,s)
29	H	Ph	3,5-diMe-Ph	—	FA:369,N1:2.07(6H,s),6.68(1H,s),13.11 (2H,s)
30	H	Ph	3-AcO-Ph	—	FA: 399
31	H	Ph	3-Br-Ph	—	FA:419,N1:6.99-7.03(1H,m), 7.73-7.75(2H,m), 13.12 (2H,s)
32	H	Ph	2,6-diF-Ph	—	FA: 377

Table 17




EX	R <sup>2</sup>	A	B	Sa	DAT
33	H	Ph	3-(MeOCO)Ph	—	FA: 399
34	MeOCO	Ph	Ph	—	FA: 399
35	H	3,4-diF-Ph	3-O <sub>2</sub> N-Ph	—	FA:422,N2:6.46-6.56(1H, m),6.83-6.93(2H,m), 7.30-7.60(5H,m),7.72-7.87(1H, m), 7.98-8.22(2H,m), 12.79 (2H,s)
36	H	Ph	3,5-diCl-Ph	—	FA:342,N1:7.28-7.33(5H,m), 7.74-7.77(2H,m),13.15 (2H, s)
37	O <sub>2</sub> N	3,5-diF-Ph	Ph	—	FA: 422
38	H	3,5-diF-Ph	Thiop-2-yl	—	FA: 383
39	H	3-(Cl-CH <sub>2</sub> )Ph	Ph	—	FA: 389
40	H	3-H <sub>2</sub> N-Ph	3,5-diF-Ph	HCl	FA:392,N1:6.90-7.24(9H, m), 7.71-7.79(2H,m), 13.10 (2H,s)
41	H	3-H <sub>2</sub> N-4-Me-Ph	3,5-diF-Ph	HCl	FA:406
42	H <sub>2</sub> N	Ph	Ph	—	FA: 356
43	H	3-H <sub>2</sub> N-Ph	Ph	—	FA: 356
44	H	Ph	3-(EtCONH)Ph	—	FA: 412
45	H	3-(MeCONH)Ph	3-(MeCONH)Ph	—	FA: 455
46	PhCONH	Ph	Ph	—	FA: 460
47	H	Ph	3-{Ph(CH <sub>2</sub> ) <sub>4</sub> CONH}Ph	—	FA: 516
48	H	3-(Py-4-yl)CH <sub>2</sub> CONHPh	3,5-diF-Ph	HCl	FA: 511
49	H	3,5-diF-Ph	3-(Et <sub>2</sub> NCH <sub>2</sub> CONH)Ph	HCl	FA: 505
50	EtCONH	Ph	Ph	—	FA: 412
51	PhCH <sub>2</sub> CONH	Ph	Ph	—	FA: 474
52	H	3,5-diF-Ph	3-{Et <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> CONH}Ph	HCl	FA: 519
79	H	Ph	3-(Py-4-yl)(CH <sub>2</sub> ) <sub>2</sub> NHCH <sub>2</sub> }Ph	HCl	FA: 475
80	H	4-ClCH <sub>2</sub> -Ph	4-(Mo-4-yl)CH <sub>2</sub> Ph	—	FA: 488
81	H	3-{Et <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> }Ph	3-{Et <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> }Ph	—	FA: 511
82	H	Ph	3-HO-Ph	—	FA:357,N1:6.45-6.48(1H,m), 9.22(1H,s), 13.04 (2H, s)
83	H	4-(4-yl)CH <sub>2</sub> NHPh	3,5-diF-Ph	HCl	FA:472,N1:4.28(2H,s),6.43-7.02(7H,m),7.25-.80(5H,m), 9.06(1H,s), 13.09(2H,s), 14.3-14.8 (2H, m)

Table 18



EX	R <sup>2</sup>	A	B	Sa	DAT
84	H	Ph	Py-3-ylCH <sub>2</sub> NHPh	HCl	FA: 447
85	H	Ph	3-(4-AcNHPhCH <sub>2</sub> NH)Ph	—	FA: 503
86	H	3-(PrNH)Ph	3,5-diF-Ph	HCl	FA:434,N1:0.93(3H,t), 1.45-1.65 (2H,m), 3.01 (2H,t),6.90-7.37(9H, m),7.70-7.80 (2H,m), 13.13 (2H,s)
87	Py-3-ylCH <sub>2</sub> NH	Ph	Ph	—	FA:447,N1:4.32(m,2H), 6.40-6.89(3H,m),6.99-7.43(12H,m),7.78-8.64 (3H, m), 12.79 (2H, m)
88	H	3,5-diF-Ph	3-(Ph(CH <sub>2</sub> ) <sub>3</sub> NH)Ph	HCl	FA:510
89	4(AcNH)PhCH <sub>2</sub> NH	Ph	Ph	—	FA: 503
90	H	3,5-diF-Ph	3-(MeO(CH <sub>2</sub> ) <sub>2</sub> NH)Ph	HCl	FA:450,N1:3.13-3.25 (2H,m),3.30(3H,s),3.42-3.50(2H,m),6.65-7.05 (7H,m),7.25-7.37(2H, m),7.66-7.78 (2H,m), 13.02-13.18 (2H,m)
117	H	3,4,5,-triF-Ph	3-O <sub>2</sub> N-Ph	—	FA:440
118	H	3,5-diF-Ph	Naph-2-yl	—	FA:427
119	H	Benzo[b]Thiop-2-yl	Benzo[b]Thiop-2-yl	—	FA:453
120	H	3,5-diF-Ph	4-Cl-3-NO <sub>2</sub> -Ph	—	FA:456
121	H	3,5-diF-Ph	3-O <sub>2</sub> N-2-Me-Ph	—	FA:436
122	H	3,5-diF-Ph	Benzo[b]Thiop-2-yl	—	FA:433
123	H	3,5-diF-Ph	4-CN-Ph	—	FA:402
124	H	3,5-diF-Ph	3-O <sub>2</sub> N-4-MeO-Ph	—	FA:452
125	H	3,5-diF-Ph	5-O <sub>2</sub> N-Fu-2-yl	—	FA:412
126	PhCO-OCH <sub>2</sub>	Ph	Ph	—	FA:475
127	Ac	3,5-diF-Ph	Ph	—	FA:419
128	3-O <sub>2</sub> N-PhCH <sub>2</sub> NH	3,5-diF-Ph	Ph	—	FA:527
129	O <sub>2</sub> N	3,5-diF-Ph	3-Me-Ph	—	FA:436
130	3-O <sub>2</sub> N-PhCONH	3,5-diF-Ph	Ph	—	FA:541
131	TBS-OCH <sub>2</sub>	3,5-diF-Ph	Ph	—	FA:521
132	PhCH <sub>2</sub> O	3,5-diF-Ph	Ph	—	FA:483
133	O <sub>2</sub> N	3,5-diF-Ph	3-O <sub>2</sub> N-Ph	—	FA:467
134	O <sub>2</sub> N	3,5-diF-Ph	3-F-Ph	—	FA:440
135	O <sub>2</sub> N	3,5-diF-Ph	3-O <sub>2</sub> N-4-Me-Ph	—	FA:481
136	PhCH <sub>2</sub> S	3,5-diF-Ph	Ph	—	FA:498
137	Py-3-ylCH <sub>2</sub> NHSO <sub>2</sub>	3,5-diF-Ph	Ph	—	FA:547
138	O <sub>2</sub> N	3,5-diF-Ph	4-Me-Ph	—	FA:436


Table 19



EX	R <sup>2</sup>	A	B	Sa	DAT
139	O <sub>2</sub> N	3,5-diF-Ph	4-F <sub>3</sub> C-O-Ph	—	FA:506
140	O <sub>2</sub> N	3,5-diF-Ph	3-F <sub>3</sub> C-Ph	—	FA:490
141	O <sub>2</sub> N	3,5-diF-Ph	2-F <sub>3</sub> C-O-Ph	—	FA:505
142	O <sub>2</sub> N	3,5-diF-Ph	4-Cl-Ph	—	FA:456
143	O <sub>2</sub> N	3,5-diF-Ph	4-F-Ph	—	FA:440
144	O <sub>2</sub> N	3,5-diF-Ph	2,3-diMe-Ph	—	FA:450
145	O <sub>2</sub> N	3,5-diF-Ph	3-MeO-Ph	—	FA:452
146	Py-3-ylCH <sub>2</sub> O	3,5-diF-Ph	2,3-diMe-Ph	—	FA:512
147	Py-3-ylCH <sub>2</sub> O	3,5-diF-Ph	3-Me-Ph	—	FA:498
148	O <sub>2</sub> N	3,5-diF-Ph	2-MeO-Ph	—	FA:452
149	O <sub>2</sub> N	3,5-diF-Ph	6-Cl-Py-3-yl	—	FA:457
150	O <sub>2</sub> N	3,5-diF-Ph	3-CN-Ph	—	FA:447
151	O <sub>2</sub> N	3,5-diF-Ph	Naph-1-yl	—	FA:472
152	O <sub>2</sub> N	3,5-diF-Ph	4-CN-Ph	—	FA:447
153	O <sub>2</sub> N	3,5-diF-Ph	3,4-diF-Ph	—	FA:458
154	O <sub>2</sub> N	3,5-diF-Ph	4-F <sub>3</sub> C-Ph	—	FA:490
155	O <sub>2</sub> N	3,5-diF-Ph	3-AcO-Ph	—	FA:480
156	O <sub>2</sub> N	3,5-diF-Ph	Thiop-2-yl	—	FA:428
157	O <sub>2</sub> N	3,5-diF-Ph	2,3-diMeO-Ph	—	FA:482
158	O <sub>2</sub> N	3-Me-Ph	3-Me-Ph	—	FA:414
160	O <sub>2</sub> N	3,5-diF-Ph	3-AcO-2-Me-Ph	—	FA:494
161	4-O <sub>2</sub> N-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:541
162	O <sub>2</sub> N	4-F-Ph	3-Me-Ph	—	FA:418
163	O <sub>2</sub> N	2-MeO-Ph	3-Me-Ph	—	FA:430
164	O <sub>2</sub> N	2,3-diMe-Ph	3-Me-Ph	—	FA:428
165	Py-3-yl(CH <sub>2</sub> ) <sub>2</sub>	3,5-diF-Ph	3-Me-Ph	—	FA:496
166	4-O <sub>2</sub> N-PhCH <sub>2</sub> O	3,5-diF-Ph	3-Me-Ph	—	FA:541
167	H	3,4,5-triF-Ph	3-H <sub>2</sub> N-Ph	HCl	FA:410
168	H	3,5-diF-Ph	3-H <sub>2</sub> N-2-Me-Ph	HCl	FA:406
169	H	3,5-diF-Ph	3-H <sub>2</sub> N-4-Cl-Ph	-	FA:426
170	H	3,5-diF-Ph	3-H <sub>2</sub> N-4-MeO-Ph	-	FA:422
171	H	3,5-diF-Ph	5-H <sub>2</sub> N-Fu-2-yl	—	FA:382




Table 20




EX	R <sup>2</sup>	A	B	Sa	DAT
159	4-O <sub>2</sub> N-PhCH <sub>2</sub> NH	3,5-diF-Ph	2,3-diMe-Ph	—	FA:555
172	3-H <sub>2</sub> N-PhCONH	3,5-diF-Ph	Ph	—	FA:511
174	3-H <sub>2</sub> N-PhCH <sub>2</sub> NH	3,5-diF-Ph	Ph	—	FA:497
175	H <sub>2</sub> N	3,5-diF-Ph	3,5-diF-Ph	—	FA:428
176	H <sub>2</sub> N	3,5-diF-Ph	3-H <sub>2</sub> N-4-Me-Ph	—	FA:421
177	H <sub>2</sub> N	3,5-diF-Ph	3-H <sub>2</sub> N-Ph	—	FA:407
178	H <sub>2</sub> N	3,5-diF-Ph	2-Me-Ph	—	FA:406
179	H <sub>2</sub> N	3,5-diF-Ph	3-F <sub>3</sub> C-O-Ph	—	FA:476
180	H <sub>2</sub> N	3,5-diF-Ph	3-F <sub>3</sub> C-Ph	—	FA:460
181	H <sub>2</sub> N	3,5-diF-Ph	4-F-Ph	—	FA:410
182	H <sub>2</sub> N	3,5-diF-Ph	2,3-diMe-Ph	—	FA:420
183	H <sub>2</sub> N	3,5-diF-Ph	3-MeO-Ph	—	FA:422
184	H <sub>2</sub> N	3,5-diF-Ph	2-MeO-Ph	—	FA:422
185	H <sub>2</sub> N	3,5-diF-Ph	2,5-diF-Ph	—	FA:428
186	H <sub>2</sub> N	3,5-diF-Ph	6-CN-Py-3-yl	—	FA:427
187	H <sub>2</sub> N	3,5-diF-Ph	3,5-diMe-Ph	—	FA:420
188	H <sub>2</sub> N	3,5-diF-Ph	3-CN-Ph	—	FA:417
189	H <sub>2</sub> N	3,5-diF-Ph	4-CN-Ph	—	FA:417
190	H <sub>2</sub> N	3,5-diF-Ph	Naph-1-yl	—	FA:442
191	H <sub>2</sub> N	3,5-diF-Ph	3,4-diF-Ph	—	FA:428
192	H <sub>2</sub> N	3,5-diF-Ph	5-Br-Py-3-yl	—	FA:501
193	H <sub>2</sub> N	3,5-diF-Ph	3-AcO-Ph	—	FA:450
194	H <sub>2</sub> N	3,5-diF-Ph	2,3-diMeO-Ph	—	FA:452
195	H <sub>2</sub> N	3,5-diF-Ph	Thiop-2-yl	—	FA:398
196	H <sub>2</sub> N	3,5-diF-Ph	Ph	—	FA:392
197	H <sub>2</sub> N	3,5-diF-Ph	3-Me-Ph	—	FA:406
198	H <sub>2</sub> N	3,5-diF-Ph	3-AcO-2-Me-Ph	—	FA:464
199	4-H <sub>3</sub> N-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:511
200	4-H <sub>2</sub> N-PhCH <sub>2</sub> O	3,5-diF-Ph	3-Me-Ph	—	FA:542
201	H <sub>2</sub> N	4-F-Ph	3-Me-Ph	—	FA:388
202	H <sub>2</sub> N	2-MeO-Ph	3-Me-Ph	—	FA:400
203	H <sub>2</sub> N	2,3-diMe-Ph	3-Me-Ph	—	FA:398
204	PhCONH	3,5-diF-Ph	Ph	—	FA:496
205	4-(Et <sub>2</sub> NCO)PhCH <sub>2</sub> NH	3,5-diF-Ph	Ph	—	FA:581
206	Py-2-ylCH <sub>2</sub> CONH	3,5-diF-Ph	Ph	—	FA:511
207	(4-MeO-Ph)(CH <sub>2</sub> ) <sub>2</sub> CONH	3,5-diF-Ph	Ph	—	FA:554
208	3-F-PhCONH	3,5-diF-Ph	Ph	—	FA:514
209	4-MeO-PhCH <sub>2</sub> CONH	3,5-diF-Ph	Ph	—	FA:540
210	4-Me <sub>2</sub> N-PhCONH	3,5-diF-Ph	Ph	—	FA:539

Table 21



EX	R <sup>2</sup>	B	Sa	DAT
211	4-Ac-PhCONH	Ph	—	FA:538
212	2-Me-PhCONH	Ph	—	FA:510,N1:2.42(3H, s), 6.90-7.00(3H,m),7.11-7.21(3H,m), 7.29-7.34(4H,m),7.38-7.40(1H,m),7.49(1H,d,J=7.8Hz), 7.56(1H, dd, J = 8.8, 1.5Hz), 7.67(1H,d,J=8.8Hz),8.37(1H, s),10.45(1H,s),13.10(1H,s),13.14 (1H, s)
213	4-AcNH-PhCONH	Ph	—	FA:553
214	Py-3-yl-CONH	Ph	—	FA:497
215	3-Cl-PhCONH	Ph	—	FA:530
216	MeOCO(CH <sub>2</sub> ) <sub>2</sub> CONH	Ph	—	FA:506
217	4-MeOCO-PhCONH	Ph	—	FA:554
218	4-Me-PhCH <sub>2</sub> CONH	Ph	—	FA:524
219	BenzoIM-5-ylCONH	Ph	—	FA:536
220	Thiop-2-ylCOCONH	Ph	—	FA:530
221	3-AcNH-PhCH <sub>2</sub> NH	Ph	—	FA:488
222	3-(cPrNHCO)PhCH <sub>2</sub> NH	Ph	—	FA:565
223	4-(4-F-PhNHNHCO)PhCH <sub>2</sub> NH	Ph	—	FA:634
224	4-H <sub>2</sub> NCO-PhCH <sub>2</sub> NH	Ph	—	FA:525
225	4-(iPrNHCO)PhCH <sub>2</sub> NH	4-F-Ph	—	FA:585
226	Pyra-2-ylCONH	3-Me-Ph	—	FA:512
227	Py-3-ylCONH	4-F-Ph	—	FA:515
228	cPrNHCO	2-MeO-Ph	—	FA:595
229	H <sub>2</sub> NCO	2-MeO-Ph	—	FA:555
230	PhO-CONH	Ph	—	FA:512
231	MeSO <sub>2</sub> NH	Ph	—	FA:470
232	4-AcNH-PhSO <sub>2</sub> NH	Ph	—	FA:589
233	4-F-PhSO <sub>2</sub> NH	3-Me-Ph	—	FA:566
234	4-MeO-PhSO <sub>2</sub> NH	4-F-Ph	—	FA:600
235	3-F <sub>3</sub> C-PhSO <sub>2</sub> NH	2-MeO-Ph	—	FA:551
236		2-MeO-Ph	—	FA:587
237		Ph	—	FA:587
238	6-F <sub>3</sub> C-Py-3-ylCH <sub>2</sub> NH	Ph	HCl	FA:550
239	Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:483
240	Me(Py-3-ylCH <sub>2</sub> )N	Ph	—	FA:497
241	4-AcNH-PhCH <sub>2</sub> NH	Ph	—	FA:539,N1:2.02(3H,s),4.22(2H,m),6.34(1H,m),6.65-7.53(15H,m),9.88(1H,s),12.81 (2H, m)
242		Ph	—	FA:639
243		Ph	HCl	FA:669

Table 22



EX	$R^2$	B	Sa	DAT
244	1-Me-5-F <sub>3</sub> C-Pyrazo-3-ylThiop-2-ylCH <sub>2</sub> NH	Ph	HCl	FA:636,N1:3.99(3H,s), 4.58(2H,s), 6.82-6.95(5H, m), 7.09-7.20(4H, m), 7.23 (1H,d,J=2.5Hz), 7.27-7.29 (2H,m), 7.40(1H,d,J=3.4Hz), 7.51(1H, d, J = 8.7Hz), 12.89(1H, s), 12.95(1H, s)
245	Py-4-ylCH <sub>2</sub> NH	Ph	—	FA:483,N1:4.34(2H,d,J=5.8Hz),6.55(1H,t,J=5.8Hz),6.68 (1H,dd,J = 8.8,2.5Hz),6.82-6.94(4H,m),7.09-7.18(3H, m), 7.29 (2H, d, J = 8.3Hz), 7.38 (2H,d,J = 5.8Hz), 7.45 (1H, d, J = 8.8Hz),8.51 (2H, d, J =5.8Hz),12.77(1H,s),12.88(1H,s)
246	(4-Me <sub>2</sub> N-PhCH <sub>2</sub> ) <sub>2</sub> N	Ph	—	FA:658
247	4-HOOC-PhCH <sub>2</sub> NH	Ph	—	FA:526
248	3-HO-5-HOCH <sub>2</sub> -2-Me-Py-4ylCH <sub>2</sub> NH	Ph	—	FA:543
249	6-Cl-imidazo[1,2-a]Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:556
250	IM-3-ylCH <sub>2</sub> NH	Ph	2HCl	FA:472
251	4-AcNH-PhCH <sub>2</sub> NH	3-Me-Ph	—	FA:553,N1:2.02(3H,m),2.14(3H,s),4.21(2H,d),6.80-7.15 (10H,m),7.25-7.55(5H,m), 9.85-9.90(1H,m),12.76-12.90(2H, m)
252	Thiaz-2-ylCH <sub>2</sub> NH	Ph	—	FA:489
253	PhCOCH <sub>2</sub> NH	Ph	—	FA:510
254	1-Oxido-Py-4-ylCH <sub>2</sub> NH	Ph	—	FA:499
255	5-(4-Cl-Ph)Fu-2-ylCH <sub>2</sub> NH	Ph	HCl	FA:582
256	Thiaz-5-ylCH <sub>2</sub> NH	Ph	2HCl	FA:489,N1:4.77(2H,s), 6.88-6.98(3H,m),7.11-7.21(4H, m), 7.33(2H,d,J=7Hz),7.55(1H, br),7.66(1H,t,J=9Hz),7.82 (1H,d,J=4 Hz),9.15(1H,d,J = 9 Hz), 13.10(2H, br)
257	5,6-diCl-Py-3-yl-CH <sub>2</sub> NH	Ph	—	FA:551
258	Pyrazi-3-ylCH <sub>2</sub> NH	Ph	—	FA:484
259	5-Br-Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:561
260	Pyrim-5-ylCH <sub>2</sub> NH	Ph	—	FA:484 N1:4.35(2H, m), 6.47 - 9.12(15H,m),12.84(2H,m)
261	6-Me-Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:497
262	2-Me-Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:497
263	3-F <sub>3</sub> C-PhCH <sub>2</sub> NH	Ph	—	FA:550
264	2-OH-PhCH <sub>2</sub> NH	Ph	—	FA:498
265	1-Me-IM-2-ylCH <sub>2</sub> NH	Ph	—	FA:486
266	3-HOOC-PhCH <sub>2</sub> NH	Ph	—	FA:526
267	3-MeO-PhCH <sub>2</sub> NH	Ph	—	FA:512
268	4-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:527
269	Py-3-ylCH <sub>2</sub> NHCH <sub>2</sub>	Ph	—	FA:497
270	4-F-PhCH <sub>2</sub> NH	Ph	—	FA:500

Continuation of Table 22



271	6-AcNH-Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:540
272	4-HO-PhCH <sub>2</sub> NH	Ph	—	FA:498
273	PhCH <sub>2</sub> NH	Ph	—	FA:482
276	4-AcNH-PhNHCH <sub>2</sub>	Ph	—	FA:539
277	Py-3-ylNHCH <sub>2</sub>	Ph	—	FA:483
278	4-AcNH-PhCH <sub>2</sub> NH	3-F-Ph	—	FA:557
279	(Py-3-yl)(CH <sub>2</sub> ) <sub>2</sub> NHCH <sub>2</sub>	Ph	—	FA:511
280	4-Br-Thiop-2-ylCH <sub>2</sub> NH	Ph	—	FA:567
281	Py-3-ylCH <sub>2</sub> NH	3-F-Ph	—	FA:501
282	2-H <sub>2</sub> N-Py-3-ylCH <sub>2</sub> NH	Ph	HCl	FA:498
283		Ph	HCl	FA:584
284	4-CN-PhCH <sub>2</sub> NH	Ph	—	FA:507
285	4-MeOCO-PhCH <sub>2</sub> NH	Ph	—	FA:540
274	4-AcNH-PhCH <sub>2</sub> NH	3-H <sub>2</sub> N-4-Me-Ph	—	FA:568, N1:1.89(3H,s), 2.02(3H,s), 4.21(2H,m), 4.77(2H,s), 6.25-6.95(9H,m), 7.25-7.55(5H,m), 9.88(1H,s), 12.68(1H,s), 12.77(1H, s)
275	4-AcNH-PhCH <sub>2</sub> NH	3-H <sub>2</sub> N-Ph	—	FA:554, N1:2.02(3H,s), 4.21(2H,m), 5.00(2H,s), 6.45-7.00 (10H, m), 7.28-7.56(5H,m), 9.88(1H,s), 12.70(1H,s), 12.79(1H, s)
286	4-AcNH-PhCH <sub>2</sub> NH	4-Me-Ph	—	FA:553

Table 23




EX	R <sup>2</sup>	B	Sa	DAT
287	4-AcNH-PhCH <sub>2</sub> NH	2-F <sub>3</sub> C-O-Ph	—	FA:622
288	4-AcNH-PhCH <sub>2</sub> NH	3-Cl-Ph	—	FA:573 N1 : 2.02(3H, s), 4.22(2H, d, J = 4.9Hz), 6.36(1H, t, J = 5.8Hz), 6.70(1H, dd, J = 8.8, 2.0Hz), 6.88-6.92(3H, m), 6.98(1H, tt, J = 8.3, 2.4Hz), 7.12-7.16(1H, m), 7.20-7.26(3H, m), 7.32(2H, d, J = 8.7Hz), 7.44(1H, d, J = 8.8Hz), 7.53(2H, d, J = 8.3Hz), 9.88(1H, s), 12.81(1H, s), 12.90(1H, s)
289	4-AcNH-PhCH <sub>2</sub> NH	4-F-Ph	—	FA:557 N1:2.01(3H, s), 4.21(2H, m), 6.34(1H, s), 6.65-7.53(14H, m), 9.87(1H, s), 12.80(2H, m)
290	4-AcNH-PhCH <sub>2</sub> NH	2,3-diMe-Ph	—	FA:567,N1:2.01(3H,s),2.02(3H,s),2.10(3H,s),4.21(2H,s),6.35(1H,br),6.67-6.73(3H,m), 6.77-6.94(5H,m),7.31(2H,d,J=9Hz), 7.43 (1H, d,J=9Hz),7.52(2H,d,J=9Hz),9.88(1H,s), 12.85 (1H,s),12.94(1H, s)
291	4-AcNH-PhCH <sub>2</sub> NH	3-MeO-Ph	—	FA:569 N1 : 2.02(3H, s), 3.66(3H, s), 4.22(2H, d, J = 5.3Hz), 6.35(1H, t, J = 5.6Hz), 6.67(1H, dd, J = 8.8, 2.5Hz), 6.71(1H, dd, J = 8.3, 2.5Hz), 6.79(1H, s), 6.86-6.87(4H, m), 6.92-6.97(1H, m), 7.02(1H, t, J = 7.8Hz), 7.31(2H, d, J = 8.3Hz), 7.42(1H, d, J = 8.8Hz), 7.53(2H, d, J = 8.3Hz), 9.89(1H, s), 12.78(1H, s), 12.87(1H, s)
292	4-HO-3-O <sub>2</sub> N-PhNHCH <sub>2</sub>	Ph	—	FA:543
293	4-AcNH-PhCH <sub>2</sub> NH	2-MeO-Ph	—	FA:569,N1:2.02(3H,s),3.63(3H,s),4.22(2H,d,J=5.3Hz),6.34(1H,t,J=5.9Hz),6.58 (1H,d,J=8.3 Hz),6.61-6.71 (2H, m),6.75-6.78(2H,m),6.87-6.92(2H,m),7.00-7.07(2H,m),7.31(2H,d, J=8.3Hz),7.42(1H,d,J=8.8Hz),7.53(2H,d,J=8.8Hz), 9.88(1H,s), 12.84(1H,s), 12.94 (1H, s)
294	4-AcNH-PhCH <sub>2</sub> NH	3,5-diMe-Ph	—	FA:567
295	4-AcNH-PhCH <sub>2</sub> NH	3-CN-Ph	—	FA:564
296	4-AcNH-PhCH <sub>2</sub> NH	2-F-Ph	—	FA:557,N2:2.19(3H, s), 4.35(2H, s), 6.46-7.87(15H, m), 9.93(1H, s), 12.65(2H, m)
297	4-AcNH-PhCH <sub>2</sub> NH	4-CN-Ph	—	FA:564,N1 :2.01(3H, s), 4.21(2H, m), 6.37(1H, s),6.67-7.87(14H,m),9.88(1H,s),12.86(2H, m)
298	4-AcNH-PhCH <sub>2</sub> NH	Naph-1-yl	—	FA:589,N1:2.02(3H, s), 4.23(2H, s), 6.38(1H, br), 6.47-6.58(3H, m), 6.71(1H, dd, J = 2 Hz, 9 Hz), 6.91(1H, s), 7.17-7.26(2H, m), 7.32(2H, d, J = 9 Hz), 7.42-7.54(5H, m), 7.64(1H, d, J = 8 Hz), 7.75(1H, d, J = 8 Hz), 8.14(1H, d, J = 8 Hz),9.89(1H,s), 12.90(1H, s), 12.99(1H, s)
299	4-AcNH-PhCH <sub>2</sub> NH	3,4-diF-Ph	—	FA:575
300	3-CN-PhCH <sub>2</sub> NH	4-F-Ph	—	FA:525
301	4-AcNH-PhCH <sub>2</sub> NH	2-Cl-Ph	—	FA:573,N1 : 2.02(3H, s), 4.22(2H, d, J = 5.3Hz), 6.38(1H, t, J = 5.9Hz), 6.70(1H, dd, J = 8.8, 2.0Hz), 6.83-6.96(4H, m), 7.05-7.17(4H, m), 7.31(2H, d, J = 8.3Hz), 7.44(1H, d, J = 8.3Hz), 7.53(2H, d, J = 8.8Hz), 9.88(1H, s), 12.87(1H, s), 12.97(1H, s)

Continuation  
of Table 23


302	4-AcNH-PhCH <sub>2</sub> NH	2,5-diF-Ph	—	FA:575,N1:2.02(3H, s), 4.22(2H, m), 6.30-6.75(2H, m), 6.82-7.10(7H, m), 7.27-7.57(5H, m), 9.88(1H, s), 12.80-13.05(2H, m)
303	4-AcNH-PhCH <sub>2</sub> NH	2-F <sub>3</sub> C-Ph	—	FA:607,N1:2.02(3H, s), 4.22(2H, m), 6.38(1H, m), 6.60-6.95(5H, m), 7.22-7.56(9H, m), 9.88(1H, s), 12.85-13.00(2H, m)
304	4-AcNH-PhCH <sub>2</sub> NH	6-Cl-Py-3-yl	—	FA:574
305	4-AcNH-PhCH <sub>2</sub> NH	5-Br-Py-3-yl	—	FA:618
306	4-AcNH-PhCH <sub>2</sub> NH	3-Br-Ph	—	FA:616
307	4-AcNH-PhCH <sub>2</sub> NH	3-AcO-Ph	—	FA:597
308	4-HO-3-MeO-PhCH <sub>2</sub> NH	3,5-diMeO-Ph	—	FA:588
309	4-HOOC-PhCH <sub>2</sub> NH	4-F-Ph	—	FA:544
310	4-AcNH-PhCH <sub>2</sub> NH	Thiop-2-yl	—	FA:545
311	4-AcNH-PhCH <sub>2</sub> NH	2,3-diMeO-Ph	—	FA:599,N1:2.02(3H,s),3.65(3H,S),3.66(3H,s),4.21(2H,d,J=5.8Hz),6.35(1H,t,J=5.9 Hz), 6.59-6.61(1H,m),6.71(1H,dd,J=2.8,8.8Hz), 6.74-6.79(4H,m),6.85-6.91(2H,m),7.32(2H,d,J=8.3Hz),7.43(1H,d,J=8.8Hz),7.53(2H,d,J=8.8Hz),9.88(1H,s),12.84(1H,s),12.93(1H, s)

Table 24



EX	R <sup>2</sup>	B	Sa	DAT
312	4-[Mo-4-yl(CH <sub>2</sub> ) <sub>2</sub> O]PhCH <sub>2</sub> NH	4-F-Ph	—	FA:629
313	4-HO-PhCH <sub>2</sub> NH	2-MeO-Ph	—	FA:528
314	3-EtO-4-MeO-PhCH <sub>2</sub> NH	3-Me-Ph	—	FA:570
315	4-(Me <sub>2</sub> N(CH <sub>2</sub> ) <sub>3</sub> O)PhCH <sub>2</sub> NH	3-Me-Ph	—	FA:597
316	1,3-benzoThiaz-6-ylCH <sub>2</sub> NH	3-Me-Ph	—	FA:553
317	4-[Mo-4-yl(CH <sub>2</sub> ) <sub>2</sub> O]PhCH <sub>2</sub> NH	2-MeO-Ph	—	FA:641
318	3-HOOC-PhCH <sub>2</sub> NH	2-MeO-Ph	—	FA:556
319	3-CN-PhCH <sub>2</sub> NH	3-Me-Ph	—	FA:521
320	1-PhCH <sub>2</sub> -Pipe-4-ylCH <sub>2</sub> NH	Ph	HCl	FA:579
321	PhCH <sub>2</sub> NH	Ph	—	FA:482
322	Naph-2-yl-CH <sub>2</sub> NH	Ph	—	FA:532
323	4-Me-PhCH <sub>2</sub> NH	Ph	—	FA:496
324	3-MeO-PhCH <sub>2</sub> NH	Ph	—	FA:512
325	2-CN-PhCH <sub>2</sub> NH	Ph	—	FA:507
326	3-F <sub>3</sub> C-PhCH <sub>2</sub> NH	Ph	—	FA:550
327	3-Br-PhCH <sub>2</sub> NH	Ph	—	FA:560
328	4-HO-PhCH <sub>2</sub> NH	Ph	—	FA:498
329	4-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:527
330	4-MeS-PhCH <sub>2</sub> NH	Ph	—	FA:528
331	2-MeO-Naph-1-yl-CH <sub>2</sub> NH	Ph	—	FA:562

Table 25






EX	R <sup>2</sup>	B	Sa	DAT
332	5,6,7,8,-tetrahydro-Naph-2-yl-CH <sub>2</sub> NH	Ph	—	FA:536
333	2,3-dihydro-benzo[b]Fu-5-ylCH <sub>2</sub> NH	Ph	—	FA:524
334		Ph	—	FA:526
335		Ph	—	FA:540
336	3,4-diMeO-PhCH <sub>2</sub> NH	Ph	—	FA:542
337	2,5-diF-PhCH <sub>2</sub> NH	Ph	—	FA:518
338	3,5-diF <sub>3</sub> C-PhCH <sub>2</sub> NH	Ph	—	FA:618
339	5-Et-Fu-2-ylCH <sub>2</sub> NH	Ph	—	FA:500
340	Thiop-3-ylCH <sub>2</sub> NH	Ph	—	FA:488
341	1-MeO-CO(CH <sub>2</sub> ) <sub>2</sub> -Pyr-2-ylCH <sub>2</sub> NH	Ph	—	FA:557
342	Pen-NH	Ph	—	FA:462
343	PhCH <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> NH	Ph	—	FA:526
344	Ph(CH <sub>2</sub> ) <sub>3</sub> NH	Ph	—	FA:510
345	Me(Py-3-yl)CHNH	Ph	—	FA:497
346	4-Pen-PhCH <sub>2</sub> NH	Ph	—	FA:552
347	biPh-4-ylCH <sub>2</sub> NH	Ph	—	FA:558
348	4-F <sub>3</sub> C-PhCH <sub>2</sub> NH	Ph	—	FA:550
349	2-Cl-PhCH <sub>2</sub> NH	Ph	—	FA:516
350	4-MeO-CO-PhCH <sub>2</sub> NH	Ph	—	FA:540
351	3-CN-PhCH <sub>2</sub> NH	Ph	—	FA:507
352	4-Me <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:525
353	4-Pyrroli-1-yl-PhCH <sub>2</sub> NH	Ph	—	FA:551
354	4-PrO-PhCH <sub>2</sub> NH	Ph	—	FA:540
355	4-HOCOCH <sub>2</sub> O-PhCH <sub>2</sub> NH	Ph	—	FA:556
356	4-PhO-PhCH <sub>2</sub> NH	Ph	—	FA:574
357	3-(3-F <sub>3</sub> C-PhO)PhCH <sub>2</sub> NH	Ph	—	FA:642
358	4-PhCH <sub>2</sub> O-PhCH <sub>2</sub> NH	Ph	—	FA:588
359	4-biPhO-PhCH <sub>2</sub> NH	Ph	—	FA:650
360	2-(4-Cl-PhS)-PhCH <sub>2</sub> NH	Ph	—	FA:624
361	6-MeO-Naph-2-ylCH <sub>2</sub> NH	Ph	—	FA:562
362	1-HO-Naph-2-ylCH <sub>2</sub> NH	Ph	—	FA:548
363	9H-Fluoren-2-ylCH <sub>2</sub> NH	Ph	—	FA:570
364	2-Cl-5-F-PhCH <sub>2</sub> NH	Ph	—	FA:534
365	3,5-diHO-PhCH <sub>2</sub> NH	Ph	—	FA:514
366	2-HO-3-MeO-PhCH <sub>2</sub> NH	Ph	—	FA:528
367	2-HO-4-Me <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:569
368	2-HO-5-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:543
369	4-HO-3-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:543
370	4-HO-3-MeO-PhCH <sub>2</sub> NH	Ph	—	FA:528
371	2,4-diMeO-PhCH <sub>2</sub> NH	Ph	—	FA:542
372	3-MeO-2-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:557
373	4-(Mo-1-yl)-2-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:612

Table 26



EX	R <sup>2</sup>	B	Sa	DAT
374	3,5-diCl-6-HO-PhCH <sub>2</sub> NH	Ph	—	FA:566
375	3,4-diMeO-2-O <sub>2</sub> N-PhCH <sub>2</sub> NH	Ph	—	FA:587
376	4-MeO-5,6-diMe-PhCH <sub>2</sub> NH	Ph	—	FA:540
377	3-HO-4,5-diMeO-PhCH <sub>2</sub> NH	Ph	—	FA:558
378	1-PhSO <sub>2</sub> -Pyr-2-ylCH <sub>2</sub> NH	Ph	—	FA:611
379	5-AcOCH <sub>2</sub> -Fu-2-ylCH <sub>2</sub> NH	Ph	—	FA:544
380	5-Me-Thiop-2-ylCH <sub>2</sub> NH	Ph	—	FA:502
381	5-Thiop-2-ylThiop-2-ylCH <sub>2</sub> NH	Ph	—	FA:570
382	4-Br-Thiop-2-ylCH <sub>2</sub> NH	Ph	—	FA:566
383	2-Ph-IM-4-ylCH <sub>2</sub> NH	Ph	—	FA:548
384	2-H <sub>2</sub> N-Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:498
385	Indol-3-ylCH <sub>2</sub> NH	Ph	—	FA:521
386	1-(4-Me-PhSO <sub>2</sub> )indol-3-ylCH <sub>2</sub> NH	Ph	—	FA:675
387	3-Me-benzo[b]Thiop-2-ylCH <sub>2</sub> NH	Ph	—	FA:552
388	quinolin-3-ylCH <sub>2</sub> NH	Ph	—	FA:533
389	5-PhCH <sub>2</sub> O-1H-pyrrolo[2,3-c]Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:628
390	PrNH	Ph	—	FA:434
391	cHex-CH <sub>2</sub> NH	Ph	—	FA:488
392	PhCH <sub>2</sub> NH	Ph	—	FA:496
393	1-Tr-BenzolM-5-ylCH <sub>2</sub> NH	Ph	—	FA:550
394	4-AcNH-PhCH <sub>2</sub> NH	3-OH-Ph	—	FA:555
395	Me(Py-3-yl)C=N	Ph	—	FA:495
396	Me(Py-3-yl)CH <sub>2</sub> NH	Ph	HCl	FA:497
397	H	6-Me-Py-3-yl	—	FA:392
398	H	1-Me-benzolM-5-yl	—	FA:431
399	Me(HO)CH <sub>2</sub>	Ph	—	FA:421
400	1-Oxido-Py-3-ylCH <sub>2</sub> NH	Ph	—	FA:499
401	HO	Ph	—	FA:393



Table 27






Ex	Str	DAT
110		FA: 391
111		FA: 394
450		FA:542,FN:540

Table 28




EX	R <sup>2</sup>	A	B	Sa	DAT
402	Py-3-ylCH <sub>2</sub> O	3,5-diF-Ph	Ph	Oxal	FA:483
403	PhCH <sub>2</sub> SO <sub>2</sub>	3,5-diF-Ph	Ph	—	FA:530
404	1-Oxido-Py-3-ylCH=N	3,5-diF-Ph	Ph	—	FA:497
405	HOCH <sub>2</sub>	3,5-diF-Ph	Ph	—	FA:407
406	BenzolM-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	Ph	—	FA:522
407	PhNHCSNH	3,5-diF-Ph	Ph	—	FA:527
408	Py-3-ylNHCONH	3,5-diF-Ph	Ph	—	FA:512
409	HCO	3,5-diF-Ph	Ph	—	FA:405
410	4-AcNHPhCH <sub>2</sub> NH	3-Me-Ph	3-Me-Ph	—	FA:531
411	H <sub>2</sub> N	3-Me-Ph	3-Me-Ph	—	FA:384
412	4-H <sub>2</sub> N-PhCH <sub>2</sub> NH	3,5-diF-Ph	2,3-diMe-Ph	—	FN:523
413	4-IPrNHOC-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:581
414	4-HOOC-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:540
415	4-[Pyrroli-1-yl(CH <sub>2</sub> ) <sub>2</sub> O]-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:609
416	PhO-CONH	3,5-diF-Ph	Ph	—	FA:512
417	4-MeOCH <sub>2</sub> CONH-PhCH <sub>2</sub> (MeOCH <sub>2</sub> CO)N	3,5-diF-Ph	2,3-diMe-Ph	—	FA:669
418	4-cBuCONH-PhCH <sub>2</sub> NH	3,5-diF-Ph	2,3-diMe-Ph	—	FA:607
419	4-Et <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> CONH-PhCH <sub>2</sub> NH	3,5-diF-Ph	2,3-diMe-Ph	Oxal	FA:652
420	4-MeNHCH <sub>2</sub> CONH-PhCH <sub>2</sub> NH	3,5-diF-Ph	2,3-diMe-Ph	Oxal	FA:596
422	1,3-Thiaz-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	Oxal	FA:503
423	2-H <sub>2</sub> N-1,3-Thiaz-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:518
424	2-AcNH-1,3-Thiaz-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:560
425	4-MeO-Ph	3,5-diF-Ph	3-Me-Ph	—	FA:497
426	1-PhCH <sub>2</sub> -Pipe-4-ylCH <sub>2</sub> NH	3,5-diF-Ph	Ph	HCl	FA:579
427	4-Me-PhSO <sub>2</sub> (Me)N	3,5-diF-Ph	Ph	—	FA:560
428	1,3-Thiaz-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	2,3-diMe-Ph	Oxal	FA:517
430	4-MeO-PhNHCO	3,5-diF-Ph	Ph	—	FA:526
431	PhCH <sub>2</sub> O-CO	3,5-diF-Ph	Ph	—	FA:511
432	HOOC	3,5-diF-Ph	Ph	—	FA:421
433	4-F-PhCO(Me)N	3,5-diF-Ph	Ph	—	FA:528
434	MeNH	3,5-diF-Ph	Ph	—	FA:406
435	2-tBuO-CONH-4-Cl-1,3-Thiaz-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:652
436	2-tBuO-CONH-4-Cl-1,3-Thiaz-5-ylCH=N	3,5-diF-Ph	3-Me-Ph	—	FA:650
437	4-AcNH-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-AcO-2-Me-Ph	—	FA:611

Table 29




EX	R <sup>2</sup>	A	B	Sa	DAT
421	1,3-Thiaz-5-ylCH <sub>2</sub> NH	3,5-diF-Ph	2-MeO-Ph	Oxal	FA:519,N1:3.64(3H,s), 4.54(2H,s),6.59(1H,d,J=8Hz),6.68-6.72(2H, m), 6.77(2H,dd, J=2Hz,8Hz), 6.90(1H,tt,J=2Hz,9Hz), 6.97(1H,d,J=2Hz),7.01-7.07(2H,m),7.45(1H,d,J=9Hz),7.89(1H,s),8.96(1H,s), 12.90(1H,s),12.96(1H, s)
438	4-AcNH-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-HO-2-Me-Ph	—	FA:569
439	4-Ac(Me)N-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:567
440	4-F <sub>3</sub> CCONH-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:607
441	4-MeSO <sub>2</sub> NH-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:589
442	4-AcNH-PhCH <sub>2</sub> O	3,5-diF-Ph	3-Me-Ph	—	FA:554,N1:2.05(3H,s), 2.51(3H,s),5.06(2H,s), 6.86-7.07(7H,m),7.14(1H,d,J=7Hz),7.39 (1H,s),7.41(2H,d,J=9Hz),7.61(2H,d,J=9Hz),7.64(1H,s),9.98(1H,s),13.06(2H,s)
443	4-AcNH-PhCH <sub>2</sub> NH	4-F-Ph	3-Me-Ph	—	FA:535
444	4-AcNH-PhCH <sub>2</sub> NH	2-MeO-Ph	3-Me-Ph	—	FA:547
445	4-AcNH-PhCH <sub>2</sub> NH	2,3-diMe-Ph	3-Me-Ph	—	FA:545
446	4-MeO-Ph(Me)NCO	3,5-diF-Ph	Ph	—	FA:540
447	4-[BocHNC(NBoc)NH]-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	—	FA:753
448	4-[H <sub>2</sub> NC(NH)NH]-PhCH <sub>2</sub> NH	3,5-diF-Ph	3-Me-Ph	HCl	FA:553
449	4-MeSO <sub>2</sub> NH-PhCH <sub>2</sub> (MeSO <sub>2</sub> )N	3,5-diF-Ph	3-Me-Ph	—	FA:667

Table 30




Ex	R <sup>1</sup> -Z <sup>1</sup>	R <sup>2</sup> -Z <sup>2</sup>	DAT
24	CH	N	FA:342,N1:7.04-7.38(11H, m),8.00(1H, m),8.34(1H, m), 13.13-13.19(2H,m)
25	N	CH	FA:342,N1:7.04-7.34(10H,m),7.66(1H,m),8.41(1H,m),8.91(1H,m),13.21(m,2H)

Table 31



Ex	R <sup>2</sup>	R <sup>6</sup>	Sa	DAT
114	H	Me	—	FA:355
115	H	PhCH <sub>2</sub>	—	FA:429
116	Me	Et <sub>2</sub> N	HCl	FA:426

Table 32





Ex	R <sup>16</sup>	DAT	Ex	R <sup>16</sup>	DAT
53	Ac	FA: 455	71	BuSO <sub>2</sub>	FA: 611
54	PhCO	FA: 579	72	Me <sub>2</sub> NSO <sub>2</sub>	FA: 585
55	2-F <sub>3</sub> C-PhCO	FA: 715	73	PhCH <sub>2</sub> SO <sub>2</sub>	FA: 679
56		FA: 973	74	PhSO <sub>2</sub>	FA: 651
57	2-Me-PhCO	FA: 607	75	3-Me-PhSO <sub>2</sub>	FA: 679
58	PhCH <sub>2</sub> CO	FA: 607	76	2,4-di-F-PhSO <sub>2</sub>	FA: 723
59	2-PyCO	FA: 581	77	4-MeO-PhSO <sub>2</sub>	FA: 711
60	MeOCH <sub>2</sub> CO	FA: 515	78	3-O <sub>2</sub> N-PhSO <sub>2</sub>	FA: 741
61	4-F-PhCO	FA: 615	91	cHex-CH <sub>2</sub>	FA: 563
62	iPrCO	FA: 511	92	PhCH <sub>2</sub>	FA: 551
63	3-Cl-PhCO	FA: 647	93	2-(EtO)PhCH <sub>2</sub>	FA: 639
64	3-MeO-PhCO	FA: 639	94	3-BrPhCH <sub>2</sub>	FA: 709
65	EtOCOCO	FA: 571	95	3-MePhCH <sub>2</sub>	FA: 579
66	4-CN-PhCO	FA: 629	96	3-NO <sub>2</sub> PhCH <sub>2</sub>	FA: 641
67	iPrSO <sub>2</sub>	FA: 583	97	4-(MeOCO)PhCH <sub>2</sub>	FA: 667
68	4-F-PhSO <sub>2</sub>	FA: 687	98	2,4-diF-PhCH <sub>2</sub>	FA: 623
69	2-F <sub>3</sub> C-PhSO <sub>2</sub>	FA: 787	99	3-PyCH <sub>2</sub>	FA: 553
70	MeSO <sub>2</sub>	FA: 527	100	4-IM-CH <sub>2</sub>	FA: 530

Table 33



EX	R <sup>1</sup>	R <sup>2</sup>	A	B	Sa	DAT
101	H	H	2-Me-1,3-Thiaz-4-yl	Ph	—	FA:362,N1:2.30(3H,s), 7.64 (1H, s), 13.04(2H, s)
102	H	MeO	Ph	Ph	—	FA: 371
103	H	H	4-Py	Ph	—	FA: 342
104	Me	H	Ph	Ph	—	FA: 355
105	H	H	3-Py	Ph	—	FA:342,N1:8.24(1H,dd, $J=$ 1.5,4.4),8.42(1H,d, $J=$ 1.5),13.18(2H, s)
106	H	H	Ph	3-{Mo-4-yl(CH <sub>2</sub> ) <sub>2</sub> O}Ph	—	FA: 470
107	H	H	Ph	3-Me <sub>2</sub> NPh	—	FA: 384
108	H	H	Ph	3-{Me(PhCH <sub>2</sub> )NCH <sub>2</sub> }Ph	HCl	FA: 474
109	H	H	3,5-diF-Ph	Py-3-yl	HCl	FA:378,N1:7.79-7.81(2H, m),8.74(1H,d, $J=$ 1.5), 13.28 (2H, s)
112	H	H	5-HO-Py-3-yl	3,5-diF-Ph	—	FA: 394
113	H	OH	3-MePh	3-MePh	—	FA: 383

### Industrial Applicability

5                Since the compounds of the invention have sex hormone-decreasing effect based on GnRH receptor antagonism, they are useful for treating sex hormone-dependent diseases, for example, prostate cancer, breast cancer, endometriosis, uterine leiomyoma, and the like (Proc. Natl. Acad. Sci. USA, 10    87, 7100-7104 (1990)).

In the following, *in vitro* GnRH receptor antagonistic effect of the pharmaceutical drugs and compounds of the invention was evaluated by inhibition of binding of <sup>125</sup>I-D-Trp<sup>6</sup>-LHRH to human GnRH receptor.

#### 15    1. Test of *in vitro* GnRH receptor antagonism

(1) Preparation of human GnRH receptor-expressing CHO (Chinese hamster ovary) cells

Expression of human GnRH was carried out in a similar manner to a common method for protein expression (Chapter 9

In: Current Protocols In Molecular Biology: ed. by F. M.

Ausubel et al., Greene Publishing Associates and Wiley-

5 Interscience, 9.0.1-9.9.6 (1987), S. S. Kaker et al., Biol.

Biophys. Res. Commun. 189, 289-295 (1992), R. Grosse et al.,

Mol. Endocrinol. 11, 1305-1318 (1997)). CHO cells were

cultured (medium:  $\alpha$ MEM, 10% FCS, containing an antibiotic-

antimycotic agent) with an expression vector which has human

10 GnRH receptor gene (SEQ ID No.: 1) and a transfection reagent

FuGENE6 (manufactured by Boehringer-Mannheim) for 24 hours for

transfection, whereby the CHO cells stably expressing human

GnRH receptor (SEQ ID No.: 2, S. S. Kaker et al., Biol.

Biophys. Res. Commun, 189, 289-295 (1992)) were obtained. The

15 expression of the aimed receptor was confirmed by PCR method.

(2) Preparation of CHO cell membrane fraction containing human

GnRH receptor

The CHO cells expressing the human GnRH receptor ( $3 \times 10^8$  cells), prepared in the above (1), were suspended in phosphate

20 buffered saline (PBS) and centrifuged for 3 minutes at 100xG.

The pellet suspended in 100 ml of a homogenate buffer (10mM

$\text{NaHCO}_3$ , 5mM EDTA (ethylenediaminetetraacetic acid), pH 7.5) was

homogenized using a Polytron homogenizer. After the resulting

homogenate was centrifuged for 15 minutes at 400xG, the

25 supernatant was centrifuged for 1 hour at 100,000xG. The

membrane fraction obtained as a pellet was suspended in 60 ml

of storage buffer (25mM Tris-HCl, 1 mM EDTA, 10  $\mu\text{g/ml}$  a

protease inhibitor (Pefabloc SC (manufactured by Merck)), 1  $\mu\text{g/ml}$  pepstatin A, 20  $\mu\text{g/ml}$  leupeptin, 0.03% sodium azide, pH 7.5), and stored at  $-80^{\circ}\text{C}$  until use.

(3) Measurement of  $^{125}\text{I}$ -D-Trp<sup>6</sup>-LHRH binding inhibition

5           The membrane fractions of CHO cells expressing the human GnRH receptor which were prepared in the above (2) were diluted with assay buffer (HBSS (Hanks balanced salt solution), 20 mM HEPES, 0.1% BSA (bovine serum albumin), 100  $\mu\text{g/ml}$  bacitracin, pH 7.4) so as to be 20  $\mu\text{g/tube}$ , and were  
10 dispensed into tubes in an amount of 148  $\mu\text{l}$ . Incubation (at  $4^{\circ}\text{C}$  for 3 hours) was initiated by adding a test compound having a different concentration dissolved in DMSO (2  $\mu\text{l}$ ) and 0.1 nM  $^{125}\text{I}$ -D-Trp<sup>6</sup>-LHRH (50  $\mu\text{l}$ ) (SEQ ID No.: 3). DMSO (2  $\mu\text{l}$ ) was added instead of the test compound for measuring a total binding and  
15 100  $\mu\text{M}$  LHRH (2  $\mu\text{l}$ ) (SEQ ID No.: 4) for non-specific binding. Incubation was terminated and bound and free ligands were separated by rapid filtration through a Whatman glass filter (GF/B) treated with 0.5% polyethyleneimine. The filters were washed three times with 1 ml of assay buffer and the  
20 radioactivity on the filter was measured using a  $\gamma$ -counter. Binding inhibition (%) (PMB) of each test compound at various concentrations were determined according to the following equation:  $\text{PMB} = (\text{SB} - \text{NSB}) / (\text{TB} - \text{NSB}) \times 100$  (wherein TB: total binding radioactivity, SB: radioactivity when a test compound  
25 was added, NSB: non-specific binding radioactivity). PMB at each concentration of each test compound was plotted and the concentration of the test compound at which PMB equals 50%

(IC<sub>50</sub> value) was determined. As a result of the test, it was confirmed that the compound No. 178a in Table 6 and the compounds of Examples 40, 43, 79, 83, 87, 132, 146, 147, 169, 209, 224, 239, 241, 245, 251, 256, 258, 290, 293, 400, 402, 421, 422 and 423 have IC<sub>50</sub> values of 10<sup>-10</sup>M to 10<sup>-9</sup>M. In particular, the compounds of the invention having a dihydrobenzimidazol-2-ylidene-substituted propane-1,3-dione skeleton have GnRH receptor binding inhibitory activities equal to that of Cetrorelix, a peptide antagonist, which is commercialized at present.

## 2. Test of *in vivo* GnRH receptor antagonism

*In vivo* GnRH antagonistic effect of each compound was evaluated by inhibitory effect on the increase of serum testosterone induced by GnRH administration in rats.

GnRH (30 ng/rat; Peptide Institute, LH-RH (human)) (SEQ ID No.: 4) was administered intramuscularly to Wistar male rats (9 wk old; Japan SLC). Each test compound was dissolved or suspended in a 6.7% DMSO, 6.7% PEG400, 6.7% Tween80 aqueous solution and was administered orally (30 mg/kg) 3 hours before the GnRH administration. Blood specimens were obtained 1 hour after the GnRH administration, and the serum concentration of testosterone was measured by a specific radioimmunoassay (IATRON Labs.).

The inhibitory activity (%) (IA) of the test compound was determined according to the following equation:  $IA = (Tc - Ts) / (Tc - Tn) \times 100$ ; wherein Tn: the serum testosterone concentration of rats to which GnRH was not administered, Tc:



the serum testosterone concentration of rats to which the solvent was administered instead of the test compound, Ts: the serum testosterone concentration of rats to which the test compound was administered (when the concentration is decreased to Tn, IA becomes 100%). As a result of the test, the compound No. 63a in Table 5, the compound No. 167a, 169a and 173a, and the compounds of Examples 40, 212, 241, 244, 245, 251, 256, 260, 274, 275, 288, 289, 290, 291, 293, 296, 297, 298, 301, 302, 303, 311 and 421 showed inhibitory activities larger than 50%.

From the test 1 and 2 in the above, it has been proved that the compounds of the invention have a sex hormone-decreasing effect based on a strong GnRH receptor antagonism, and hence are useful for treating sex hormone-dependent diseases, for example, prostate cancer, breast cancer, endometriosis, uterine leiomyoma, and the like (C. Huggins & C. V. Hodges, Cancer Res. 1, 293-297 (1941), L. Bokser et al., Proc. Natl. Acad. Sci. USA, 87, 7100-7104 (1990)).

1/4

SEQUENCE LISTING

<110> Yamanouchi pharm. co., ltd.

<120> Propane derivatives

<130> GnRH

<140>

<141>

<150> JP2000-204425

<151> 2000-07-05

<150> JP2001-153372

<151> 2001-05-23

<160> 4

<170> PatentIn Ver. 2.1

<210> 1

<211> 987

<212> DNA

<213> Homo sapiens

<300>

<303> Biol. Biophys. Res. Comm.

<304> 189

<306> 289-295

<307> 1992

<400> 1

```
atggcaaaca gtgcctctcc tgaacagaat caaaatcact gttcagccat caacaacagc 60
atcccactga tgcaggggcaa cctccccact ctgaccttgt ctggaaagat ccgagtgaac 120
gttactttct tctttttct gctctctgcg acctttaatg cttctttctt gttgaaactt 180
cagaagtgga cacagaagaa agagaaaagg aaaaagctct caagaatgaa gctgctctta 240
aaacatctga ccttagccaa cctgttggag actctgatig tcatgccact ggatgggatg 300
```

2/4

tggaacatta cagtccaatg gtatgctgga gagttactct gcaaagttct cagttatcta 360  
 aagcttttct ccatgtatgc ccagcccttc atgatgggtg tgatcagcct ggaccgctcc 420  
 ctggctatca cgaggccctt agctttgaaa agcaacagca aagtcggaca gtccatgggtt 480  
 ggccctggcct ggatcctcag tagtgtcttt gcaggaccac agttatacat cttcaggatg 540  
 attcatctag cagacagctc tggacagaca aaagtittct ctcaatgtgt aacacactgc 600  
 agtttttcac aatgggtggc tcaagcattt tataactttt tcaccttcag ctgcctcttc 660  
 atcatccctc ttttcatcat gctgatctgc aatgcaaaaa tcattctcac cctgacacgg 720  
 gtccttcac aggaccccca cgaactacaa ctgaatcagt ccaagaacaa tataccaaga 780  
 gcacggctga agactctaaa aatgacgggt gcatttgcca cttcatttac tgtctgctgg 840  
 actccctact atgtcctagg aatttgggtat tggtttgatc ctgaaatgtt aaacaggttg 900  
 tcagaccag taaatcactt cttctttctc ttgctttt taaacccatg ctttgatcca 960  
 cttatctatg gatatttttc tctgtga 987

&lt;210&gt; 2

&lt;211&gt; 328

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

Met Ala Asn Ser Ala Ser Pro Glu Gln Asn Gln Asn His Cys Ser Ala

1

5

10

15

Ile Asn Asn Ser Ile Pro Leu Met Gln Gly Asn Leu Pro Thr Leu Thr

20

25

30

Leu Ser Gly Lys Ile Arg Val Thr Val Thr Phe Phe Leu Phe Leu Leu

35

40

45

Ser Ala Thr Phe Asn Ala Ser Phe Leu Leu Lys Leu Gln Lys Trp Thr

50

55

60

Gln Lys Lys Glu Lys Gly Lys Lys Leu Ser Arg Met Lys Leu Leu Leu

65

70

75

80

Lys His Leu Thr Leu Ala Asn Leu Leu Glu Thr Leu Ile Val Met Pro

85

90

95

3/4

Leu Asp Gly Met Trp Asn Ile Thr Val Gln Trp Tyr Ala Gly Glu Leu  
 100 105 110

Leu Cys Lys Val Leu Ser Tyr Leu Lys Leu Phe Ser Met Tyr Ala Pro  
 115 120 125

Ala Phe Met Met Val Val Ile Ser Leu Asp Arg Ser Leu Ala Ile Thr  
 130 135 140

Arg Pro Leu Ala Leu Lys Ser Asn Ser Lys Val Gly Gln Ser Met Val  
 145 150 155 160

Gly Leu Ala Trp Ile Leu Ser Ser Val Phe Ala Gly Pro Gln Leu Tyr  
 165 170 175

Ile Phe Arg Met Ile His Leu Ala Asp Ser Ser Gly Gln Thr Lys Val  
 180 185 190

Phe Ser Gln Cys Val Thr His Cys Ser Phe Ser Gln Trp Trp His Gln  
 195 200 205

Ala Phe Tyr Asn Phe Phe Thr Phe Ser Cys Leu Phe Ile Ile Pro Leu  
 210 215 220

Phe Ile Met Leu Ile Cys Asn Ala Lys Ile Ile Phe Thr Leu Thr Arg  
 225 230 235 240

Val Leu His Gln Asp Pro His Glu Leu Gln Leu Asn Gln Ser Lys Asn  
 245 250 255

Asn Ile Pro Arg Ala Arg Leu Lys Thr Leu Lys Met Thr Val Ala Phe  
 260 265 270

Ala Thr Ser Phe Thr Val Cys Trp Thr Pro Tyr Tyr Val Leu Gly Ile  
 275 280 285

Trp Tyr Trp Phe Asp Pro Glu Met Leu Asn Arg Leu Ser Asp Pro Val  
 290 295 300

4/4

Asn His Phe Phe Phe Leu Phe Ala Phe Leu Asn Pro Cys Phe Asp Pro  
305 310 315 320

Leu Ile Tyr Gly Tyr Phe Ser Leu  
325

&lt;210&gt; 3

&lt;211&gt; 10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:labeled Tyr  
with 125I and substituted with D form of Trp

&lt;400&gt; 3

Glu His Trp Ser Tyr Trp Leu Arg Pro Gly  
1 5 10

&lt;210&gt; 4

&lt;211&gt; 10

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

Glu His Trp Ser Tyr Gly Leu Arg Pro Gly  
1 5 10

1/4

## SEQUENCE LISTING

<110> Yamanouchi pharm. co., ltd.

<120> Propane derivatives

<130> GnRH

<140>

<141>

<150> JP2000-204425

<151> 2000-07-05

<150> JP2001-153372

<151> 2001-05-23

<160> 4

<170> PatentIn Ver. 2.1

<210> 1

<211> 987

<212> DNA

<213> Homo sapiens

<300>

<303> Biol. Biophys. Res. Comm.

<304> 189

<306> 289-295

<307> 1992

<400> 1

atggcaaaca gtgcctctcc tgaacagaat caaaatcact gttcagccat caacaacagc 60  
atcccaactga tgcagggcaa cctccccact ctgaccttgt ctggaaagat ccgagtgaac 120  
gttactttct tctttttct gctctctgag acctttaatg cttctttctt gttgaaactt 180  
cagaagtgga cacagaagaa agagaaaggg aaaaagctct caagaatgaa gctgctotta 240  
aaacatctga ccttagccaa cctgttgag actctgattg tcatgccact ggatgggatg 300

2/4

```

tggaacatta cagtcgaatg gtatgctgga gagttactct gcaaagttct cagttatcta 360
aagcttttct ccattgatgc ccagccttc atgatggtgg tgatcagcct ggaccgctcc 420
ctggctatca cgaggccctt agctttgaaa agcaacagca aagtcggaca gtccatggtt 480
ggcctggcct ggatcctcag tagtgtcttt gcaggaccac agttatacat cttcaggatg 540
attcatctag cagacagctc tggacagaca aaagttttct ctcaatgtgt aacacactgc 600
agtttttcac aatggtggca tcaagcattt tataactttt tcaccttcag ctgcctcttc 660
atcatccctc ttttcatcat gctgatctgc aatgcaaaaa tcactttcac cctgacacgg 720
gtccttcac aggaccccca cgaactacaa ctgaatcagt ccaagaacaa tataccaaga 780
gcacggctga agactctaaa aatgacgggt gcatttgcca cttcatttac tgtctgctgg 840
actccctact atgtcctagg aatttggtat tggtttgatc ctgaaatgtt aaacagggtg 900
tcagaccag taaatcactt cttctttctc ttgctttt taaacccatg ctttgatcca 960
cttatctatg gatatttttc tctgtga 987

```

&lt;210&gt; 2

&lt;211&gt; 328

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

Met Ala Asn Ser Ala Ser Pro Glu Gln Asn Gln Asn His Cys Ser Ala
  1                      5                      10                     15

```

```

Ile Asn Asn Ser Ile Pro Leu Met Gln Gly Asn Leu Pro Thr Leu Thr
          20                      25                     30

```

```

Leu Ser Gly Lys Ile Arg Val Thr Val Thr Phe Phe Leu Phe Leu Leu
          35                      40                     45

```

```

Ser Ala Thr Phe Asn Ala Ser Phe Leu Leu Lys Leu Gln Lys Trp Thr
          50                      55                     60

```

```

Gln Lys Lys Glu Lys Gly Lys Lys Leu Ser Arg Met Lys Leu Leu Leu
          65                      70                     75                     80

```

```

Lys His Leu Thr Leu Ala Asn Leu Leu Glu Thr Leu Ile Val Met Pro
          85                      90                     95

```

3/4

Leu Asp Gly Met Trp Asn Ile Thr Val Gln Trp Tyr Ala Gly Glu Leu  
 100 105 110

Leu Cys Lys Val Leu Ser Tyr Leu Lys Leu Phe Ser Met Tyr Ala Pro  
 115 120 125

Ala Phe Met Met Val Val Ile Ser Leu Asp Arg Ser Leu Ala Ile Thr  
 130 135 140

Arg Pro Leu Ala Leu Lys Ser Asn Ser Lys Val Gly Gln Ser Met Val  
 145 150 155 160

Gly Leu Ala Trp Ile Leu Ser Ser Val Phe Ala Gly Pro Gln Leu Tyr  
 165 170 175

Ile Phe Arg Met Ile His Leu Ala Asp Ser Ser Gly Gln Thr Lys Val  
 180 185 190

Phe Ser Gln Cys Val Thr His Cys Ser Phe Ser Gln Trp Trp His Gln  
 195 200 205

Ala Phe Tyr Asn Phe Phe Thr Phe Ser Cys Leu Phe Ile Ile Pro Leu  
 210 215 220

Phe Ile Met Leu Ile Cys Asn Ala Lys Ile Ile Phe Thr Leu Thr Arg  
 225 230 235 240

Val Leu His Gln Asp Pro His Glu Leu Gln Leu Asn Gln Ser Lys Asn  
 245 250 255

Asn Ile Pro Arg Ala Arg Leu Lys Thr Leu Lys Met Thr Val Ala Phe  
 260 265 270

Ala Thr Ser Phe Thr Val Cys Trp Thr Pro Tyr Tyr Val Leu Gly Ile  
 275 280 285

Trp Tyr Trp Phe Asp Pro Glu Met Leu Asn Arg Leu Ser Asp Pro Val  
 290 295 300



4/4

Asn His Phe Phe Phe Leu Phe Ala Phe Leu Asn Pro Cys Phe Asp Pro  
305 310 315 320

Leu Ile Tyr Gly Tyr Phe Ser Leu  
325

&lt;210&gt; 3

&lt;211&gt; 10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:labeled Tyr  
with 125I and substituted with D form of Trp

&lt;400&gt; 3

Glu His Trp Ser Tyr Trp Leu Arg Pro Gly  
1 5 10

&lt;210&gt; 4

&lt;211&gt; 10

&lt;212&gt; PRT


&lt;213&gt; Homo sapiens

&lt;400&gt; 4

Glu His Trp Ser Tyr Gly Leu Arg Pro Gly  
1 5 10

## Claims

1. A pharmaceutical composition comprising a propane-1,3-dione derivative represented by the general formula (I):



(R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup>: the same or different, H, NO<sub>2</sub>, CN, Halo, a hydrocarbon group which may be substituted, a heterocycle which may be substituted, a hydroxy which may be substituted, a carboxy which may be substituted, an acyl-O- which may be substituted, an acyl which may be substituted, a substituent -S(O)<sub>n<sub>101</sub></sub>- (n<sub>101</sub>: an integer of 0 to 2, the same shall apply hereinafter), H-S(O)<sub>n<sub>101</sub></sub>-, a carbamoyl which may be substituted, a sulfamoyl which may be substituted, or an amino which may be substituted, and two adjacent groups selected from the group of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> may be combined to form an aryl or a cycloalkenyl;

R<sup>5</sup> and R<sup>6</sup>: the same or different, H, Halo, a hydrocarbon group which may be substituted or an amino which may be substituted;

X<sup>1</sup> and X<sup>2</sup>: the same or different, N, S or O atom;

A and B: the same or different, an aryl which may be substituted or a heterocycle which may be substituted;

Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup> and Z<sup>4</sup>: C or N;

provided that 1) when  $X^1$  and  $X^2$  each is S or O, one or both of the corresponding  $R^5$  and  $R^6$  are absent; 2) when one to four of  $Z^1$ ,  $Z^2$ ,  $Z^3$  and/or  $Z^4$  are N, the corresponding  $R^1$ ,  $R^2$ ,  $R^3$  and/or  $R^4$  are absent)








































5 or a pharmaceutically acceptable salt thereof as the active ingredient.

2. The pharmaceutical composition according to claim 1, which is a gonadotropin releasing hormone receptor  
10 antagonist.

3. The pharmaceutical composition comprising a propane-1,3-dione derivative or a pharmaceutically acceptable salt thereof as the active ingredient according to claim 1 or  
15 2, wherein at least any one of  $X^1$  and  $X^2$  is N.

4. A propane-1,3-dione derivative or a pharmaceutically acceptable salt thereof according to claim 1, provided that the compounds 1 to 39 shown in the following  
20 table are excluded, wherein the symbol Ph means phenyl, Me means methyl, Et means ethyl, or tBu means tert-butyl.

Table 1

1. 	14. 	27. 
2. 	15. 	28. 
3. 	16. 	29. 
4. 	17. 	30. 
5. 	18. 	31. 
6. 	19. 	32. 
7. 	20. 	33. 
8. 	21. 	34. 
9. 	22. 	35. 
10. 	23. 	36. 
11. 	24. 	37. 
12. 	25. 	38. 
13. 	26. 	39. 

5. The propane-1,3-dione derivative or a pharmaceutically acceptable salt thereof according to claim 4, wherein at least any one of  $X^1$  and  $X^2$  is N.

5 6. The propane-1,3-dione derivative or a pharmaceutically acceptable salt thereof according to claim 4 or 5, wherein  $X^1$  and  $X^2$  are N at the same time.